

# BEST AVAILABLE COPY

WO 01/79468

PCT/US01/11869

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number  
**WO 01/79468 A2**

(51) International Patent Classification<sup>7</sup>: **C12N 9/00**

(21) International Application Number: PCT/US01/11869

(22) International Filing Date: 12 April 2001 (12.04.2001)

(25) Filing Language: English

(26) Publication Language: English

**(30) Priority Data:**

60/197,590	13 April 2000 (13.04.2000)	US
60/198,403	19 April 2000 (19.04.2000)	US
60/200,185	28 April 2000 (28.04.2000)	US
60/202,234	5 May 2000 (05.05.2000)	US
60/203,509	11 May 2000 (11.05.2000)	US

**(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:**

US	60/198,403 (CIP)
Filed on	19 April 2000 (19.04.2000)
US	60/197,590 (CIP)
Filed on	13 April 2000 (13.04.2000)
US	60/200,185 (CIP)
Filed on	28 April 2000 (28.04.2000)
US	60/202,234 (CIP)
Filed on	5 May 2000 (05.05.2000)
US	60/203,509 (CIP)
Filed on	11 May 2000 (11.05.2000)

(71) Applicant (for all designated States except US): **INCYTE GENOMICS, INC.** [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **POLICKY, Jennifer, L.** [US/US]; 1511 Jarvis Court, San Jose, CA 95118 (US). **HAFALIA, April** [US/US]; 2227 Calle de Primavera, Santa Clara, CA 95054 (US). **BURFORD, Neil** [GB/US]; 105 Wildwood Circle, Durham, CT 06422 (US). **RING, Huijun, Z.** [US/US]; 625 Orange Avenue, Los Altos, CA 94022 (US). **LAL, Preeti** [IN/US]; P.O. Box 5142, Santa Clara, CA 95056 (US). **TRIBOULEY, Catherine, M.** [FR/US]; 1121 Tennessee Street, #5, San Francisco, CA 94107 (US). **YAO, Monique, G.** [US/US];

111 Frederick Court, Mountain View, CA 94043 (US). **YUE, Henry** [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). **TANG, Y., Tom** [US/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). **PATTERSON, Chandra** [US/US]; 490 Sherwood Way #1, Menlo Park, CA 94025 (US). **DAS, Debopriya** [IN/US]; Apartment 3, 1179 Bonita Avenue, Mountain View, CA 94040 (US). **SANJANWALA, Madhu, S.** [US/US]; 210 Sylvia Court, Los Altos, CA 94024 (US). **GANDHI, Ameena, R.** [US/US]; 837 Roble Avenue, #1, Menlo Park, CA 94025 (US). **REDDY, Roopa** [US/US]; 1233 W. McKinley Avenue, #3, Sunnyvale, CA 94086 (US). **KHAN, Farrah, A.** [IN/US]; 333 Escuela Avenue, #221, Mountain View, CA 94040 (US). **BAUGHN, Mariah, R.** [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). **RAMKUMAR, Jaya** [IN/US]; 34359 Maybird Circle, Fremont, CA 94555 (US). **GRIFFIN, Jennifer, A.** [US/US]; 181 Irene Court, #11, Belmont, CA 94002 (US). **AU-YOUNG, Janice** [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US).

(74) Agents: **HAMLET-COX, Diana et al.**; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **DRUG METABOLIZING ENZYMES**

(57) Abstract: The invention provides human drug metabolizing enzymes (DME) and polynucleotides which identify and encode DME. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with aberrant expression of DME.

WO 01/79468 A2

## DRUG METABOLIZING ENZYMES

### TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of drug metabolizing enzymes and to the use of these sequences in the diagnosis, treatment, and prevention of autoimmune/inflammatory, cell proliferative, developmental, endocrine, eye, metabolic, and gastrointestinal disorders, including liver disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of drug metabolizing enzymes.

### BACKGROUND OF THE INVENTION

The metabolism of a drug and its movement through the body (pharmacokinetics) are important in determining its effects, toxicity, and interactions with other drugs. The three processes governing pharmacokinetics are the absorption of the drug, distribution to various tissues, and elimination of drug metabolites. These processes are intimately coupled to drug metabolism, since a variety of metabolic modifications alter most of the physicochemical and pharmacological properties of drugs, including solubility, binding to receptors, and excretion rates. The metabolic pathways which modify drugs also accept a variety of naturally occurring substrates such as steroids, fatty acids, prostaglandins, leukotrienes, and vitamins. The enzymes in these pathways are therefore important sites of biochemical and pharmacological interaction between natural compounds, drugs, carcinogens, mutagens, and xenobiotics.

It has long been appreciated that inherited differences in drug metabolism lead to drastically different levels of drug efficacy and toxicity among individuals. For drugs with narrow therapeutic indices, or drugs which require bioactivation (such as codeine), these polymorphisms can be critical. Moreover, promising new drugs are frequently eliminated in clinical trials based on toxicities which may only affect a segment of the patient group. Advances in pharmacogenomics research, of which drug metabolizing enzymes constitute an important part, are promising to expand the tools and information that can be brought to bear on questions of drug efficacy and toxicity (See Evans, W.E. and R.V. Relling (1999) Science 286:487-491).

Drug metabolic reactions are categorized as Phase I, which functionalize the drug molecule and prepare it for further metabolism, and Phase II, which are conjugative. In general, Phase I reaction products are partially or fully inactive, and Phase II reaction products are the chief excreted species. However, Phase I reaction products are sometimes more active than the original administered drugs; this metabolic activation principle is exploited by pro-drugs (e.g. L-dopa). Additionally, some nontoxic compounds (e.g. aflatoxin, benzo[a]pyrene) are metabolized to toxic

intermediates through these pathways. Phase I reactions are usually rate-limiting in drug metabolism. Prior exposure to the compound, or other compounds, can induce the expression of Phase I enzymes however, and thereby increase substrate flux through the metabolic pathways. (See Klaassen, C.D., Amdur, M.O. and J. Doull (1996) Casarett and Doull's Toxicology: The Basic Science of Poisons, McGraw-Hill, New York, NY, pp. 113-186; B.G. Katzung (1995) Basic and Clinical Pharmacology, Appleton and Lange, Norwalk, CT, pp. 48-59; G.G. Gibson and P. Skett (1994) Introduction to Drug Metabolism, Blackie Academic and Professional, London.)

Drug metabolizing enzymes (DMEs) have broad substrate specificities. This can be contrasted to the immune system, where a large and diverse population of antibodies are highly specific for their antigens. The ability of DMEs to metabolize a wide variety of molecules creates the potential for drug interactions at the level of metabolism. For example, the induction of a DME by one compound may affect the metabolism of another compound by the enzyme.

DMEs have been classified according to the type of reaction they catalyze and the cofactors involved. The major classes of Phase I enzymes include, but are not limited to, cytochrome P450 and flavin-containing monooxygenase. Other enzyme classes involved in Phase I-type catalytic cycles and reactions include, but are not limited to, NADPH cytochrome P450 reductase (CPR), the microsomal cytochrome b5/NADH cytochrome b5 reductase system, the ferredoxin/ferredoxin reductase redox pair, aldo/keto reductases, and alcohol dehydrogenases. The major classes of Phase II enzymes include, but are not limited to, UDP glucuronyltransferase, sulfotransferase, glutathione S-transferase, N-acyltransferase, and N-acetyl transferase.

#### Cytochrome P450 and P450 catalytic cycle-associated enzymes

Members of the cytochrome P450 superfamily of enzymes catalyze the oxidative metabolism of a variety of substrates, including natural compounds such as steroids, fatty acids, prostaglandins, leukotrienes, and vitamins, as well as drugs, carcinogens, mutagens, and xenobiotics. Cytochromes P450, also known as P450 heme-thiolate proteins, usually act as terminal oxidases in multi-component electron transfer chains, called P450-containing monooxygenase systems. Specific reactions catalyzed include hydroxylation, epoxidation, N-oxidation, sulfoxidation, N-, S-, and O-dealkylations, desulfation, deamination, and reduction of azo, nitro, and N-oxide groups. These reactions are involved in steroidogenesis of glucocorticoids, cortisols, estrogens, and androgens in animals; insecticide resistance in insects; herbicide resistance and flower coloring in plants; and environmental bioremediation by microorganisms. Cytochrome P450 actions on drugs, carcinogens, mutagens, and xenobiotics can result in detoxification or in conversion of the substance to a more toxic product. Cytochromes P450 are abundant in the liver, but also occur in other tissues; the enzymes are located in microsomes. (See ExPASy ENZYME EC 1.14.14.1; Prosite PDOC00081 Cytochrome P450 cysteine heme-iron ligand signature; PRINTS EP450I E-Class P450 Group I

signature; Graham-Lorence, S. and Peterson, J.A. (1996) FASEB J. 10:206-214.)

Four hundred cytochromes P450 have been identified in diverse organisms including bacteria, fungi, plants, and animals (Graham-Lorence, supra). The B-class is found in prokaryotes and fungi, while the E-class is found in bacteria, plants, insects, vertebrates, and mammals. Five subclasses or groups are found within the larger family of E-class cytochromes P450 (PRINTS EP450I E-Class P450 Group I signature).

All cytochromes P450 use a heme cofactor and share structural attributes. Most cytochromes P450 are 400 to 530 amino acids in length. The secondary structure of the enzyme is about 70% alpha-helical and about 22% beta-sheet. The region around the heme-binding site in the C-terminal part of the protein is conserved among cytochromes P450. A ten amino acid signature sequence in this heme-iron ligand region has been identified which includes a conserved cysteine involved in binding the heme iron in the fifth coordination site. In eukaryotic cytochromes P450, a membrane-spanning region is usually found in the first 15-20 amino acids of the protein, generally consisting of approximately 15 hydrophobic residues followed by a positively charged residue. (See Prosite PDOC00081, supra; Graham-Lorence, supra.)

Cytochrome P450 enzymes are involved in cell proliferation and development. The enzymes have roles in chemical mutagenesis and carcinogenesis by metabolizing chemicals to reactive intermediates that form adducts with DNA (Nebert, D.W. and Gonzalez, F.J. (1987) Ann. Rev. Biochem. 56:945-993). These adducts can cause nucleotide changes and DNA rearrangements that lead to oncogenesis. Cytochrome P450 expression in liver and other tissues is induced by xenobiotics such as polycyclic aromatic hydrocarbons, peroxisomal proliferators, phenobarbital, and the glucocorticoid dexamethasone (Dogra, S.C. et al. (1998) Clin. Exp. Pharmacol. Physiol. 25:1-9). A cytochrome P450 protein may participate in eye development as mutations in the P450 gene CYP1B1 cause primary congenital glaucoma (Online Mendelian Inheritance in Man (OMIM) \*601771 Cytochrome P450, subfamily I (dioxin-inducible), polypeptide 1; CYP1B1).

Cytochromes P450 are associated with inflammation and infection. Hepatic cytochrome P450 activities are profoundly affected by various infections and inflammatory stimuli, some of which are suppressed and some induced (Morgan, E.T. (1997) Drug Metab. Rev. 29:1129-1188). Effects observed in vivo can be mimicked by proinflammatory cytokines and interferons.

Autoantibodies to two cytochrome P450 proteins were found in patients with autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), a polyglandular autoimmune syndrome (OMIM \*240300 Autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy).

Mutations in cytochromes P450 have been linked to metabolic disorders, including congenital adrenal hyperplasia, the most common adrenal disorder of infancy and childhood; pseudovitamin D-deficiency rickets; cerebrotendinous xanthomatosis, a lipid storage disease characterized by



progressive neurologic dysfunction, premature atherosclerosis, and cataracts; and an inherited resistance to the anticoagulant drugs coumarin and warfarin (Isselbacher, K.J. et al. (1994) Harrison's Principles of Internal Medicine, McGraw-Hill, Inc. New York, NY, pp. 1968-1970; Takeyama, K. et al. (1997) *Science* 277:1827-1830; Kitataka, S. et al. (1998) *N. Engl. J. Med.* 338:653-661; OMIM \*213700 Cerebrotendinous xanthomatosis; and OMIM #122700 Coumarin resistance). Extremely high levels of expression of the cytochrome P450 protein aromatase were found in a fibrolamellar hepatocellular carcinoma from a boy with severe gynecomastia (feminization) (Agarwal, V.R. (1998) *J. Clin. Endocrinol. Metab.* 83:1797-1800).

The cytochrome P450 catalytic cycle is completed through reduction of cytochrome P450 by NADPH cytochrome P450 reductase (CPR). Another microsomal electron transport system consisting of cytochrome b5 and NADPH cytochrome b5 reductase has been widely viewed as a minor contributor of electrons to the cytochrome P450 catalytic cycle. However, a recent report by Lamb, D.C. et al. (1999; *FEBS Lett.* 462:283-288) identifies a Candida albicans cytochrome P450 (CYP51) which can be efficiently reduced and supported by the microsomal cytochrome b5/NADPH cytochrome b5 reductase system. Therefore, there are likely many cytochromes P450 which are supported by this alternative electron donor system.

Cytochrome b5 reductase is also responsible for the reduction of oxidized hemoglobin (methemoglobin, or ferrihemoglobin, which is unable to carry oxygen) to the active hemoglobin (ferrohemoglobin) in red blood cells. Methemoglobinemia results when there is a high level of oxidant drugs or an abnormal hemoglobin (hemoglobin M) which is not efficiently reduced. Methemoglobinemia can also result from a hereditary deficiency in red cell cytochrome b5 reductase (Reviewed in Mansour, A. and Lurie, A.A. (1993) *Am. J. Hematol.* 42:7-12).

Members of the cytochrome P450 family are also closely associated with vitamin D synthesis and catabolism. Vitamin D exists as two biologically equivalent prohormones, ergocalciferol (vitamin D<sub>2</sub>), produced in plant tissues, and cholecalciferol (vitamin D<sub>3</sub>), produced in animal tissues. The latter form, cholecalciferol, is formed upon the exposure of 7-dehydrocholesterol to near ultraviolet light (i.e., 290-310 nm), normally resulting from even minimal periods of skin exposure to sunlight (reviewed in Miller, W.L. and Portale, A.A. (2000) *Trends Endocrinol. Metab.* 11:315-319).

Both prohormone forms are further metabolized in the liver to 25-hydroxyvitamin D (25(OH)D) by the enzyme 25-hydroxylase. 25(OH)D is the most abundant precursor form of vitamin D which must be further metabolized in the kidney to the active form, 1 $\alpha$ ,25-dihydroxyvitamin D (1 $\alpha$ ,25(OH)<sub>2</sub>D), by the enzyme 25-hydroxyvitamin D 1 $\alpha$ -hydroxylase (1 $\alpha$ -hydroxylase). Regulation of 1 $\alpha$ ,25(OH)<sub>2</sub>D production is primarily at this final step in the synthetic pathway. The activity of 1 $\alpha$ -hydroxylase depends upon several physiological factors including the circulating level of the enzyme product (1 $\alpha$ ,25(OH)<sub>2</sub>D) and the levels of parathyroid hormone (PTH), calcitonin, insulin,

calcium, phosphorus, growth hormone, and prolactin. Furthermore, extrarenal  $1\alpha$ -hydroxylase activity has been reported, suggesting that tissue-specific, local regulation of  $1\alpha,25(\text{OH})_2\text{D}$  production may also be biologically important. The catalysis of  $1\alpha,25(\text{OH})_2\text{D}$  to 24,25-dihydroxyvitamin D ( $24,25(\text{OH})_2\text{D}$ ), involving the enzyme 25-hydroxyvitamin D

5 24-hydroxylase (24-hydroxylase), also occurs in the kidney. 24-hydroxylase can also use  $25(\text{OH})\text{D}$  as a substrate (Shinki, T. et al. (1997) Proc. Natl. Acad. Sci. U.S.A. 94:12920-12925; Miller, W.L. and Portale, A.A. supra; and references within).

Vitamin D 25-hydroxylase,  $1\alpha$ -hydroxylase, and 24-hydroxylase are all NADPH-dependent, type I (mitochondrial) cytochrome P450 enzymes that show a high degree of homology with other  
10 members of the family. Vitamin D 25-hydroxylase also shows a broad substrate specificity and may also perform 26-hydroxylation of bile acid intermediates and 25, 26, and 27-hydroxylation of cholesterol (Dilworth, F.J. et al. (1995) J. Biol. Chem. 270:16766-16774; Miller, W.L. and Portale, A.A. supra; and references within).

The active form of vitamin D ( $1\alpha,25(\text{OH})_2\text{D}$ ) is involved in calcium and phosphate  
15 homeostasis and promotes the differentiation of myeloid and skin cells. Vitamin D deficiency resulting from deficiencies in the enzymes involved in vitamin D metabolism (e.g.,  $1\alpha$ -hydroxylase) causes hypocalcemia, hypophosphatemia, and vitamin D-dependent (sensitive) rickets, a disease characterized by loss of bone density and distinctive clinical features, including bandy or bow  
20 leggedness accompanied by a waddling gait. Deficiencies in vitamin D 25-hydroxylase cause cerebrotendinous xanthomatosis, a lipid-storage disease characterized by the deposition of cholesterol and cholestanol in the Achilles' tendons, brain, lungs, and many other tissues. The disease presents with progressive neurologic dysfunction, including postpubescent cerebellar ataxia, atherosclerosis, and cataracts. Vitamin D 25-hydroxylase deficiency does not result in rickets, suggesting the  
25 existence of alternative pathways for the synthesis of  $25(\text{OH})\text{D}$  (Griffin, J.E. and Zerwekh, J.E. (1983) J. Clin. Invest. 72:1190-1199; Gamblin, G.T. et al. (1985) J. Clin. Invest. 75:954-960; and W.L. and Portale, A.A. supra).

Ferredoxin and ferredoxin reductase are electron transport accessory proteins which support at least one human cytochrome P450 species, cytochrome P450c27 encoded by the CYP27 gene (Dilworth, F.J. et al. (1996) Biochem. J. 320:267-71). A Streptomyces griseus cytochrome P450,  
30 CYP104D1, was heterologously expressed in E. coli and found to be reduced by the endogenous ferredoxin and ferredoxin reductase enzymes (Taylor, M. et al. (1999) Biochem. Biophys. Res. Commun. 263:838-42), suggesting that many cytochrome P450 species may be supported by the ferredoxin/ferredoxin reductase pair. Ferredoxin reductase has also been found in a model drug  
35 metabolism system to reduce actinomycin D, an antitumor antibiotic, to a reactive free radical species (Flitter, W.D. and Mason, R.P. (1988) Arch. Biochem. Biophys. 267:632-639).

### Flavin-containing monooxygenase (FMO)

Flavin-containing monooxygenases oxidize the nucleophilic nitrogen, sulfur, and phosphorus heteroatom of an exceptional range of substrates. Like cytochromes P450, FMOs are microsomal and use NADPH and O<sub>2</sub>; there is also a great deal of substrate overlap with cytochromes P450. The tissue distribution of FMOs includes liver, kidney, and lung.

There are five different known isoforms of FMO in mammals (FMO1, FMO2, FMO3, FMO4, and FMO5), which are expressed in a tissue-specific manner. The isoforms differ in their substrate specificities and other properties such as inhibition by various compounds and stereospecificity of reaction. FMOs have a 13 amino acid signature sequence, the components of which span the N-terminal two-thirds of the sequences and include the FAD binding region and the FATGY motif which has been found in many N-hydroxylating enzymes (Stehr, M. et al. (1998) Trends Biochem. Sci. 23:56-57; PRINTS FMOXYGENASE Flavin-containing monooxygenase signature).

Specific reactions include oxidation of nucleophilic tertiary amines to N-oxides, secondary amines to hydroxylamines and nitrones, primary amines to hydroxylamines and oximes, and sulfur-containing compounds and phosphines to S- and P-oxides. Hydrazines, iodides, selenides, and boron-containing compounds are also substrates. Although FMOs appear similar to cytochromes P450 in their chemistry, they can generally be distinguished from cytochromes P450 *in vitro* based on, for example, the higher heat lability of FMOs and the nonionic detergent sensitivity of cytochromes P450; however, use of these properties in identification is complicated by further variation among FMO isoforms with respect to thermal stability and detergent sensitivity.

FMOs play important roles in the metabolism of several drugs and xenobiotics. FMO (FMO3 in liver) is predominantly responsible for metabolizing (S)-nicotine to (S)-nicotine N-1'-oxide, which is excreted in urine. FMO is also involved in S-oxygenation of cimetidine, an H<sub>2</sub>-antagonist widely used for the treatment of gastric ulcers. Liver-expressed forms of FMO are not under the same regulatory control as cytochrome P450. In rats, for example, phenobarbital treatment leads to the induction of cytochrome P450, but the repression of FMO1.

Endogenous substrates of FMO include cysteamine, which is oxidized to the disulfide, cystamine, and trimethylamine (TMA), which is metabolized to trimethylamine N-oxide. TMA smells like rotting fish, and mutations in the FMO3 isoform lead to large amounts of the malodorous free amine being excreted in sweat, urine, and breath. These symptoms have led to the designation fish-odor syndrome (OMIM 602079 Trimethylaminuria).

### Lysyl oxidase:

Lysyl oxidase (lysine 6-oxidase, LO) is a copper-dependent amine oxidase involved in the formation of connective tissue matrices by crosslinking collagen and elastin. LO is secreted as a N-glycosylated precursor protein of approximately 50 kDa and cleaved to the mature form of the

enzyme by a metalloprotease, although the precursor form is also active. The copper atom in LO is involved in the transport of electron to and from oxygen to facilitate the oxidative deamination of lysine residues in these extracellular matrix proteins. While the coordination of copper is essential to LO activity, insufficient dietary intake of copper does not influence the expression of the apoenzyme.

5 However, the absence of the functional LO is linked to the skeletal and vascular tissue disorders that are associated with dietary copper deficiency. LO is also inhibited by a variety of semicarbazides, hydrazines, and amino nitrites, as well as heparin. Beta-aminopropionitrile is a commonly used inhibitor. LO activity is increased in response to ozone, cadmium, and elevated levels of hormones released in response to local tissue trauma, such as transforming growth factor-beta, platelet-derived  
10 growth factor, angiotensin II, and fibroblast growth factor. Abnormalities in LO activity has been linked to Menkes syndrome and occipital horn syndrome. Cytosolic forms of the enzyme have been implicated in abnormal cell proliferation (reviewed in Rucker, R.B. et al. (1998) Am. J. Clin. Nutr. 67:996S-1002S and Smith-Mungo, L.I. and Kagan, H.M. (1998) Matrix Biol. 16:387-398).

#### Dihydrofolate reductases

15 Dihydrofolate reductases (DHFR) are ubiquitous enzymes that catalyze the NADPH-dependent reduction of dihydrofolate to tetrahydrofolate, an essential step in the de novo synthesis of glycine and purines as well as the conversion of deoxyuridine monophosphate (dUMP) to deoxythymidine monophosphate (dTMP). The basic reaction is as follows:



The enzymes can be inhibited by a number of dihydrofolate analogs, including trimethoprim and methotrexate. Since an abundance of TMP is required for DNA synthesis, rapidly dividing cells require the activity of DHFR. The replication of DNA viruses (i.e., herpesvirus) also requires high  
25 levels of DHFR activity. As a result, drugs that target DHFR have been used for cancer chemotherapy and to inhibit DNA virus replication. (For similar reasons, thymidylate synthetases are also target enzymes.) Drugs that inhibit DHFR are preferentially cytotoxic for rapidly dividing cells (or DNA virus-infected cells) but have no specificity, resulting in the indiscriminate destruction of dividing cells. Furthermore, cancer cells may become resistant to drugs such as methotrexate as a  
30 result of acquired transport defects or the duplication of one or more DHFR genes (Stryer, L. (1988) Biochemistry. W.H Freeman and Co., Inc. New York. pp. 511-5619).

#### Aldo/keto reductases

Aldo/keto reductases are monomeric NADPH-dependent oxidoreductases with broad substrate specificities (Bohren, K.M. et al. (1989) J. Biol. Chem. 264:9547-9551). These enzymes  
35 catalyze the reduction of carbonyl-containing compounds, including carbonyl-containing sugars and

aromatic compounds, to the corresponding alcohols. Therefore, a variety of carbonyl-containing drugs and xenobiotics are likely metabolized by enzymes of this class.

One known reaction catalyzed by a family member, aldose reductase, is the reduction of glucose to sorbitol, which is then further metabolized to fructose by sorbitol dehydrogenase. Under normal conditions, the reduction of glucose to sorbitol is a minor pathway. In hyperglycemic states, however, the accumulation of sorbitol is implicated in the development of diabetic complications (OMIM \*103880 Aldo-keto reductase family 1, member B1). Members of this enzyme family are also highly expressed in some liver cancers (Cao, D. et al. (1998) J. Biol. Chem. 273:11429-11435).

#### Alcohol dehydrogenases

Alcohol dehydrogenases (ADHs) oxidize simple alcohols to the corresponding aldehydes. ADH is a cytosolic enzyme, prefers the cofactor NAD<sup>+</sup>, and also binds zinc ion. Liver contains the highest levels of ADH, with lower levels in kidney, lung, and the gastric mucosa.

Known ADH isoforms are dimeric proteins composed of 40 kDa subunits. There are five known gene loci which encode these subunits (a, b, g, p, c), and some of the loci have characterized allelic variants (b<sub>1</sub>, b<sub>2</sub>, b<sub>3</sub>, g<sub>1</sub>, g<sub>2</sub>). The subunits can form homodimers and heterodimers; the subunit composition determines the specific properties of the active enzyme. The holoenzymes have therefore been categorized as Class I (subunit compositions aa, ab, ag, bg, gg), Class II (pp), and Class III (cc). Class I ADH isozymes oxidize ethanol and other small aliphatic alcohols, and are inhibited by pyrazole. Class II isozymes prefer longer chain aliphatic and aromatic alcohols, are unable to oxidize methanol, and are not inhibited by pyrazole. Class III isozymes prefer even longer chain aliphatic alcohols (five carbons and longer) and aromatic alcohols, and are not inhibited by pyrazole.

The short-chain alcohol dehydrogenases include a number of related enzymes with a variety of substrate specificities. Included in this group are the mammalian enzymes D-beta-hydroxybutyrate dehydrogenase, (R)-3-hydroxybutyrate dehydrogenase, 15-hydroxyprostaglandin dehydrogenase, NADPH-dependent carbonyl reductase, corticosteroid 11-beta-dehydrogenase, and estradiol 17-beta-dehydrogenase, as well as the bacterial enzymes acetoacetyl-CoA reductase, glucose 1-dehydrogenase, 3-beta-hydroxysteroid dehydrogenase, 20-beta-hydroxysteroid dehydrogenase, ribitol dehydrogenase, 3-oxoacyl reductase, 2,3-dihydro-2,3-dihydroxybenzoate dehydrogenase, sorbitol-6-phosphate 2-dehydrogenase, 7-alpha-hydroxysteroid dehydrogenase, cis-1,2-dihydroxy-3,4-cyclohexadiene-1-carboxylate dehydrogenase, cis-toluene dihydrodiol dehydrogenase, cis-benzene glycol dehydrogenase, biphenyl-2,3-dihydro-2,3-diol dehydrogenase, N-acylmannosamine 1-dehydrogenase, and 2-deoxy-D-gluconate 3-dehydrogenase (Krozowski, Z. (1994) J. Steroid Biochem. Mol. Biol. 51:125-130; Krozowski, Z. (1992) Mol. Cell Endocrinol. 84:C25-31; and Marks, A.R. et al. (1992) J. Biol. Chem. 267:15459-15463).

### UDP glucuronyltransferase

Members of the UDP glucuronyltransferase family (UGTs) catalyze the transfer of a glucuronic acid group from the cofactor uridine diphosphate-glucuronic acid (UDP-glucuronic acid) to a substrate. The transfer is generally to a nucleophilic heteroatom (O, N, or S). Substrates include xenobiotics which have been functionalized by Phase I reactions, as well as endogenous compounds such as bilirubin, steroid hormones, and thyroid hormones. Products of glucuronidation are excreted in urine if the molecular weight of the substrate is less than about 250 g/mol, whereas larger glucuronidated substrates are excreted in bile.

UGTs are located in the microsomes of liver, kidney, intestine, skin, brain, spleen, and nasal mucosa, where they are on the same side of the endoplasmic reticulum membrane as cytochrome P450 enzymes and flavin-containing monooxygenases, and therefore are ideally located to access products of Phase I drug metabolism. UGTs have a C-terminal membrane-spanning domain which anchors them in the endoplasmic reticulum membrane, and a conserved signature domain of about 50 amino acid residues in their C terminal section (Prosite PDOC00359 UDP-glycosyltransferase signature).

UGTs involved in drug metabolism are encoded by two gene families, UGT1 and UGT2. Members of the UGT1 family result from alternative splicing of a single gene locus, which has a variable substrate binding domain and constant region involved in cofactor binding and membrane insertion. Members of the UGT2 family are encoded by separate gene loci, and are divided into two families, UGT2A and UGT2B. The 2A subfamily is expressed in olfactory epithelium, and the 2B subfamily is expressed in liver microsomes. Mutations in UGT genes are associated with hyperbilirubinemia (OMIM #143500 Hyperbilirubinemia I); Crigler-Najjar syndrome, characterized by intense hyperbilirubinemia from birth (OMIM #218800 Crigler-Najjar syndrome); and a milder form of hyperbilirubinemia termed Gilbert's disease (OMIM \*191740 UGT1).

### Sulfotransferase

Sulfate conjugation occurs on many of the same substrates which undergo O-glucuronidation to produce a highly water-soluble sulfuric acid ester. Sulfotransferases (ST) catalyze this reaction by transferring  $\text{SO}_3^-$  from the cofactor 3'-phosphoadenosine-5'-phosphosulfate (PAPS) to the substrate. ST substrates are predominantly phenols and aliphatic alcohols, but also include aromatic amines and aliphatic amines, which are conjugated to produce the corresponding sulfamates. The products of these reactions are excreted mainly in urine.

STs are found in a wide range of tissues, including liver, kidney, intestinal tract, lung, platelets; and brain. The enzymes are generally cytosolic, and multiple forms are often co-expressed. For example, there are more than a dozen forms of ST in rat liver cytosol. These biochemically characterized STs fall into five classes based on their substrate preference: arylsulfotransferase,

alcohol sulfotransferase, estrogen sulfotransferase, tyrosine ester sulfotransferase, and bile salt sulfotransferase.

ST enzyme activity varies greatly with sex and age in rats. The combined effects of developmental cues and sex-related hormones are thought to lead to these differences in ST expression profiles, as well as the profiles of other DMEs such as cytochromes P450. Notably, the high expression of STs in cats partially compensates for their low level of UDP glucuronyltransferase activity.

Several forms of ST have been purified from human liver cytosol and cloned. There are two phenol sulfotransferases with different thermal stabilities and substrate preferences. The thermostable enzyme catalyzes the sulfation of phenols such as para-nitrophenol, minoxidil, and acetaminophen; the thermolabile enzyme prefers monoamine substrates such as dopamine, epinephrine, and levadopa. Other cloned STs include an estrogen sulfotransferase and an N-acetylglucosamine-6-O-sulfotransferase. This last enzyme is illustrative of the other major role of STs in cellular biochemistry, the modification of carbohydrate structures that may be important in cellular differentiation and maturation of proteoglycans. Indeed, an inherited defect in a sulfotransferase has been implicated in macular corneal dystrophy, a disorder characterized by a failure to synthesize mature keratan sulfate proteoglycans (Nakazawa, K. et al. (1984) J. Biol. Chem. 259:13751-13757; OMIM \*217800 Macular dystrophy, corneal).

#### Galactosyltransferases

Galactosyltransferases are a subset of glycosyltransferases that transfer galactose (Gal) to the terminal N-acetylglucosamine (GlcNAc) oligosaccharide chains that are part of glycoproteins or glycolipids that are free in solution (Kolbinger, F. et al. (1998) J. Biol. Chem. 273:433-440; Amado, M. et al. (1999) Biochim. Biophys. Acta 1473:35-53). Galactosyltransferases have been detected on the cell surface and as soluble extracellular proteins, in addition to being present in the Golgi.  $\beta$ 1,3-galactosyltransferases form Type I carbohydrate chains with Gal ( $\beta$ 1-3)GlcNAc linkages. Known human and mouse  $\beta$ 1,3-galactosyltransferases appear to have a short cytosolic domain, a single transmembrane domain, and a catalytic domain with eight conserved regions. (Kolbinger, F., supra and Hennet, T. et al. (1998) J. Biol. Chem. 273:58-65). In mouse UDP-galactose: $\beta$ -N-acetylglucosamine  $\beta$ 1,3-galactosyltransferase-I region 1 is located at amino acid residues 78-83, region 2 is located at amino acid residues 93-102, region 3 is located at amino acid residues 116-119, region 4 is located at amino acid residues 147-158, region 5 is located at amino acid residues 172-183, region 6 is located at amino acid residues 203-206, region 7 is located at amino acid residues 236-246, and region 8 is located at amino acid residues 264-275. A variant of a sequence found within mouse UDP-galactose: $\beta$ -N-acetylglucosamine  $\beta$ 1,3-galactosyltransferase-I region 8 is also found in bacterial galactosyltransferases, suggesting that this sequence defines a galactosyltransferase

sequence motif (Hennet, T. supra). Recent work suggests that brainiac protein is a  $\beta$ 1,3-galactosyltransferase (Yuan, Y. et al. (1997) Cell 88:9-11; and Hennet, T. supra).

UDP-Gal:GlcNAc-1,4-galactosyltransferase (-1,4-GalT) (Sato, T. et al., (1997) EMBO J. 16:1850-1857) catalyzes the formation of Type II carbohydrate chains with Gal ( $\beta$ 1-4)GlcNAc linkages. As is the case with the  $\beta$ 1,3-galactosyltransferase, a soluble form of the enzyme is formed by cleavage of the membrane-bound form. Amino acids conserved among  $\beta$ 1,4-galactosyltransferases include two cysteines linked through a disulfide-bonded and a putative UDP-galactose-binding site in the catalytic domain (Yadav, S. and Brew, K. (1990) J. Biol. Chem. 265:14163-14169; Yadav, S.P. and Brew, K. (1991) J. Biol. Chem. 266:698-703; and Shaper, N.L. et al. (1997) J. Biol. Chem. 272:31389-31399).  $\beta$ 1,4-galactosyltransferases have several specialized roles in addition to synthesizing carbohydrate chains on glycoproteins or glycolipids. In mammals a  $\beta$ 1,4-galactosyltransferase, as part of a heterodimer with  $\alpha$ -lactalbumin, functions in lactating mammary gland lactose production. A  $\beta$ 1,4-galactosyltransferase on the surface of sperm functions as a receptor that specifically recognizes the egg. Cell surface  $\beta$ 1,4-galactosyltransferases also function in cell adhesion, cell/basal lamina interaction, and normal and metastatic cell migration. (Shur, B. (1993) Curr. Opin. Cell Biol. 5:854-863; and Shaper, J. (1995) Adv. Exp. Med. Biol. 376:95-104).

#### Glutathione S-transferase

The basic reaction catalyzed by glutathione S-transferases (GST) is the conjugation of an electrophile with reduced glutathione (GSH). GSTs are homodimeric or heterodimeric proteins localized mainly in the cytosol, but some level of activity is present in microsomes as well. The major isozymes share common structural and catalytic properties; in humans they have been classified into four major classes, Alpha, Mu, Pi, and Theta. The two largest classes, Alpha and Mu, are identified by their respective protein isoelectric points; pI ~ 7.5-9.0 (Alpha), and pI ~ 6.6 (Mu). Each GST possesses a common binding site for GSH and a variable hydrophobic binding site. The hydrophobic binding site in each isozyme is specific for particular electrophilic substrates. Specific amino acid residues within GSTs have been identified as important for these binding sites and for catalytic activity. Residues Q67, T68, D101, E104, and R131 are important for the binding of GSH (Lee, H.-C. et al. (1995) J. Biol. Chem. 270:99-109). Residues R13, R20, and R69 are important for the catalytic activity of GST (Stenberg, G. et al. (1991) Biochem. J. 274:549-555).

In most cases, GSTs perform the beneficial function of deactivation and detoxification of potentially mutagenic and carcinogenic chemicals. However, in some cases their action is detrimental and results in activation of chemicals with consequent mutagenic and carcinogenic effects. Some forms of rat and human GSTs are reliable preneoplastic markers that aid in the detection of carcinogenesis. Expression of human GSTs in bacterial strains, such as Salmonella typhimurium used in the well-known Ames test for mutagenicity, has helped to establish the role of these enzymes



in mutagenesis. Dihalomethanes, which produce liver tumors in mice, are believed to be activated by GST. This view is supported by the finding that dihalomethanes are more mutagenic in bacterial cells expressing human GST than in untransfected cells (Thier, R. et al. (1993) Proc. Natl. Acad. Sci. USA 90:8567-8580). The mutagenicity of ethylene dibromide and ethylene dichloride is increased in  
5 bacterial cells expressing the human Alpha GST, A1-1, while the mutagenicity of aflatoxin B1 is substantially reduced by enhancing the expression of GST (Simula, T.P. et al. (1993) Carcinogenesis 14:1371-1376). Thus, control of GST activity may be useful in the control of mutagenesis and carcinogenesis.

GST has been implicated in the acquired resistance of many cancers to drug treatment, the,  
10 phenomenon known as multi-drug resistance (MDR). MDR occurs when a cancer patient is treated with a cytotoxic drug such as cyclophosphamide and subsequently becomes resistant to this drug and to a variety of other cytotoxic agents as well. Increased GST levels are associated with some of these drug resistant cancers, and it is believed that this increase occurs in response to the drug agent which is then deactivated by the GST catalyzed GSH conjugation reaction. The increased GST levels then  
15 protect the cancer cells from other cytotoxic agents which bind to GST. Increased levels of A1-1 in tumors has been linked to drug resistance induced by cyclophosphamide treatment (Dirven H.A. et al. (1994) Cancer Res. 54: 6215-6220). Thus control of GST activity in cancerous tissues may be useful in treating MDR in cancer patients.

#### Gamma-glutamyl transpeptidase

20 Gamma-glutamyl transpeptidases are ubiquitously expressed enzymes that initiate extracellular glutathione (GSH) breakdown by cleaving gamma-glutamyl amide bonds. The breakdown of GSH provides cells with a regional cysteine pool for biosynthetic pathways. Gamma-glutamyl transpeptidases also contribute to cellular antioxidant defenses and expression is induced by oxidative stress. The cell surface-localized glycoproteins are expressed at high levels in  
25 cancer cells. Studies have suggested that the high level of gamma-glutamyl transpeptidases activity present on the surface of cancer cells could be exploited to activate precursor drugs, resulting in high local concentrations of anti-cancer therapeutic agents (Hanigan, M.H. (1998) Chem. Biol. Interact. 111-112:333-42; Taniguchi, N. and Ikeda, Y. (1998) Adv. Enzymol. Relat. Areas Mol. Biol. 72:239-78; Chikhi, N. et al. (1999) Comp. Biochem. Physiol. B. Biochem. Mol. Biol. 122:367-380).

#### Acyltransferase

30 N-acyltransferase enzymes catalyze the transfer of an amino acid conjugate to an activated carboxylic group. Endogenous compounds and xenobiotics are activated by acyl-CoA synthetases in the cytosol, microsomes, and mitochondria. The acyl-CoA intermediates are then conjugated with an amino acid (typically glycine, glutamine, or taurine, but also ornithine, arginine, histidine, serine,  
35 aspartic acid, and several dipeptides) by N-acyltransferases in the cytosol or mitochondria to form a

metabolite with an amide bond. This reaction is complementary to O-glucuronidation, but amino acid conjugation does not produce the reactive and toxic metabolites which often result from glucuronidation.

One well-characterized enzyme of this class is the bile acid-CoA:amino acid N-acyltransferase (BAT) responsible for generating the bile acid conjugates which serve as detergents in the gastrointestinal tract (Falany, C.N. et al. (1994) J. Biol. Chem. 269:19375-19379; Johnson, M.R. et al. (1991) J. Biol. Chem. 266:10227-10233). BAT is also useful as a predictive indicator for prognosis of hepatocellular carcinoma patients after partial hepatectomy (Furutani, M. et al. (1996) Hepatology 24:1441-1445).

#### 10 Acetyltransferases

Acetyltransferases have been extensively studied for their role in histone acetylation. Histone acetylation results in the relaxing of the chromatin structure in eukaryotic cells, allowing transcription factors to gain access to promoter elements of the DNA templates in the affected region of the genome (or the genome in general). In contrast, histone deacetylation results in a reduction in transcription by closing the chromatin structure and limiting access of transcription factors. To this end, a common means of stimulating cell transcription is the use of chemical agents that inhibit the deacetylation of histones (e.g., sodium butyrate), resulting in a global (albeit artifactual) increase in gene expression. The modulation of gene expression by acetylation also results from the acetylation of other proteins, including but not limited to, p53, GATA-1, MyoD, ACTR, TFIIE, TFIIIF and the high mobility group proteins (HMG). In the case of p53, acetylation results in increased DNA binding, leading to the stimulation of transcription of genes regulated by p53. The prototypic histone acetylase (HAT) is Gcn5 from Saccharomyces cerevisiae. Gcn5 is a member of a family of acetylases that includes Tetrahymena p55, human Gcn5, and human p300/CBP. Histone acetylation is reviewed in (Cheung, W.L. et al. (2000) Curr. Opin. Cell Biol. 12:326-333 and Berger, S.L (1999) Curr. Opin. Cell Biol. 11:336-341). Some acetyltransferase enzymes possess the alpha/beta hydrolase fold (Center of Applied Molecular Engineering Inst. of Chemistry and Biochemistry - University of Salzburg, <http://predict.sanger.ac.uk/irbm-course97/Docs/ms/>) common to several other major classes of enzymes, including but not limited to, acetylcholinesterases and carboxylesterases (Structural Classification of Proteins, <http://scop.mrc-lmb.cam.ac.uk/scop/index.html>).

#### 30 N-acetyltransferase

Aromatic amines and hydrazine-containing compounds are subject to N-acetylation by the N-acetyltransferase enzymes of liver and other tissues. Some xenobiotics can be O-acetylated to some extent by the same enzymes. N-acetyltransferases are cytosolic enzymes which utilize the cofactor acetyl-coenzyme A (acetyl-CoA) to transfer the acetyl group in a two step process. In the first step, the acetyl group is transferred from acetyl-CoA to an active site cysteine residue; in the second step,

the acetyl group is transferred to the substrate amino group and the enzyme is regenerated.

In contrast to most other DME classes, there are a limited number of known N-acetyltransferases. In humans, there are two highly similar enzymes, NAT1 and NAT2; mice appear to have a third form of the enzyme, NAT3. The human forms of N-acetyltransferase have independent regulation (NAT1 is widely-expressed, whereas NAT2 is in liver and gut only) and overlapping substrate preferences. Both enzymes appear to accept most substrates to some extent, but NAT1 does prefer some substrates (para-aminobenzoic acid, para-aminosalicylic acid, sulfamethoxazole, and sulfanilamide), while NAT2 prefers others (isoniazid, hydralazine, procainamide, dapson, aminogluthethimide, and sulfamethazine).

Clinical observations of patients taking the antituberculosis drug isoniazid in the 1950s led to the description of fast and slow acetylators of the compound. These phenotypes were shown subsequently to be due to mutations in the NAT2 gene which affected enzyme activity or stability. The slow isoniazid acetylator phenotype is very prevalent in Middle Eastern populations (approx. 70%), and is less prevalent in Caucasian (approx. 50%) and Asian (<25%) populations. More recently, functional polymorphism in NAT1 has been detected, with approximately 8% of the population tested showing a slow acetylator phenotype (Butcher, N. J. et al. (1998) Pharmacogenetics 8:67-72). Since NAT1 can activate some known aromatic amine carcinogens, polymorphism in the widely-expressed NAT1 enzyme may be important in determining cancer risk (OMIM \*108345 N-acetyltransferase 1).

#### Aminotransferases

Aminotransferases comprise a family of pyridoxal 5'-phosphate (PLP) -dependent enzymes that catalyze transformations of amino acids. Aspartate aminotransferase (AspAT) is the most extensively studied PLP-containing enzyme. It catalyzes the reversible transamination of dicarboxylic L-amino acids, aspartate and glutamate, and the corresponding 2-oxo acids, oxalacetate and 2-oxoglutarate. Other members of the family included pyruvate aminotransferase, branched-chain amino acid aminotransferase, tyrosine aminotransferase, aromatic aminotransferase, alanine:glyoxylate aminotransferase (AGT), and kynurenine aminotransferase (Vacca, R.A. et al. (1997) J. Biol. Chem. 272:21932-21937).

Primary hyperoxaluria type-1 is an autosomal recessive disorder resulting in a deficiency in the liver-specific peroxisomal enzyme, alanine:glyoxylate aminotransferase-1. The phenotype of the disorder is a deficiency in glyoxylate metabolism. In the absence of AGT, glyoxylate is oxidized to oxalate rather than being transaminated to glycine. The result is the deposition of insoluble calcium oxalate in the kidneys and urinary tract, ultimately causing renal failure (Lumb, M.J. et al. (1999) J. Biol. Chem. 274:20587-20596).

Kynurenine aminotransferase catalyzes the irreversible transamination of the L-tryptophan

metabolite L-kynurenine to form kynurenic acid. The enzyme may also catalyzes the reversible transamination reaction between L-2-aminoadipate and 2-oxoglutarate to produce 2-oxoadipate and L-glutamate. Kynurenic acid is a putative modulator of glutamatergic neurotransmission, thus a deficiency in kynurenine aminotransferase may be associated with pleiotropic effects (Buchli, R. et al. (1995) J. Biol. Chem. 270:29330-29335).

#### Catechol-O-methyltransferase

Catechol-O-methyltransferase (COMT) catalyzes the transfer of the methyl group of S-adenosyl-L-methionine (AdoMet; SAM) donor to one of the hydroxyl groups of the catechol substrate (e.g., L-dopa, dopamine, or DBA). Methylation of the 3'-hydroxyl group is favored over methylation of the 4'-hydroxyl group and the membrane bound isoform of COMT is more regiospecific than the soluble form. Translation of the soluble form of the enzyme results from utilization of an internal start codon in a full-length mRNA (1.5 kb) or from the translation of a shorter mRNA (1.3 kb), transcribed from an internal promoter. The proposed  $S_N2$ -like methylation reaction requires  $Mg^{++}$  and is inhibited by  $Ca^{++}$ . The binding of the donor and substrate to COMT occurs sequentially. AdoMet first binds COMT in a  $Mg^{++}$ -independent manner, followed by the binding of  $Mg^{++}$  and the binding of the catechol substrate.

The amount of COMT in tissues is relatively high compared to the amount of activity normally required, thus inhibition is problematic. Nonetheless, inhibitors have been developed for in vitro use (e.g., gallates, tropolone, U-0521, and 3',4'-dihydroxy-2-methyl-propionphenetropolone) and for clinical use (e.g., nitrocatechol-based compounds and tolcapone). Administration of these inhibitors results in the increased half-life of L-dopa and the consequent formation of dopamine. Inhibition of COMT is also likely to increase the half-life of various other catechol-structure compounds, including but not limited to epinephrine/norepinephrine, isoprenaline, rimiterol, dobutamine, fenoldopam, apomorphine, and  $\alpha$ -methyldopa. A deficiency in norepinephrine has been linked to clinical depression, hence the use of COMT inhibitors could be useful in the treatment of depression. COMT inhibitors are generally well tolerated with minimal side effects and are ultimately metabolized in the liver with only minor accumulation of metabolites in the body (Männistö, P.T. and Kaakkola, S. (1999) Pharmacol. Rev. 51:593-628).

#### Copper-zinc superoxide dismutases

Copper-zinc superoxide dismutases are compact homodimeric metalloenzymes involved in cellular defenses against oxidative damage. The enzymes contain one atom of zinc and one atom of copper per subunit and catalyze the dismutation of superoxide anions into  $O_2$  and  $H_2O_2$ . The rate of dismutation is diffusion-limited and consequently enhanced by the presence of favorable electrostatic interactions between the substrate and enzyme active site. Examples of this class of enzyme have been identified in the cytoplasm of all the eukaryotic cells as well as in the periplasm of several

bacterial species. Copper-zinc superoxide dismutases are robust enzymes that are highly resistant to proteolytic digestion and denaturing by urea and SDS. In addition to the compact structure of the enzymes, the presence of the metal ions and intrasubunit disulfide bonds is believed to be responsible for enzyme stability. The enzymes undergo reversible denaturation at temperatures as high as 70°C

5 (Battistoni, A. et al. (1998) J. Biol. Chem. 273:5655-5661).

Overexpression of superoxide dismutase has been implicated in enhancing freezing tolerance of transgenic Alfalfa as well as providing resistance to environmental toxins such as the diphenyl ether herbicide, acifluorfen (McKersie, B.D. et al. (1993) Plant Physiol. 103:1155-1163). In addition, yeast cells become more resistant to freeze-thaw damage following exposure to hydrogen peroxide, which causes the yeast cells to adapt to further peroxide stress by upregulating expression of superoxide dismutases. In this study, mutations to yeast superoxide dismutase genes had a more detrimental effect on freeze-thaw resistance than mutations which affected the regulation of glutathione metabolism, long suspected of being important in determining an organisms survival through the process of cryopreservation (Jong-In Park, J.-I. et al. (1998) J. Biol. Chem.

15 273:22921-22928).

Expression of superoxide dismutase is also associated with Mycobacterium tuberculosis, the organism that causes tuberculosis. Superoxide dismutase is one of the ten major proteins secreted by M. tuberculosis and its expression is upregulated approximately 5-fold in response to oxidative stress. M. tuberculosis expresses almost two orders of magnitude more superoxide dismutase than the nonpathogenic mycobacterium M. smegmatis, and secretes a much higher proportion of the expressed enzyme. The result is the secretion of ~350-fold more enzyme by M. tuberculosis than M. smegmatis, providing substantial resistance to oxidative stress (Harth, G. and Horwitz, M.A. (1999) J. Biol. Chem. 274:4281-4292).

The reduced expression of copper-zinc superoxide dismutases, as well as other enzymes with anti-oxidant capabilities, has been implicated in the early stages of cancer. The expression of copper-zinc superoxide dismutases has been shown to be lower in prostatic intraepithelial neoplasia and prostate carcinomas, compared to normal prostate tissue (Bostwick, D.G. (2000) Cancer 89:123-134).

#### Phosphodiesterases

Phosphodiesterases make up a class of enzymes which catalyze the hydrolysis of one of the two ester bonds in a phosphodiester compound. Phosphodiesterases are therefore crucial to a variety of cellular processes. Phosphodiesterases include DNA and RNA endonucleases and exonucleases, which are essential for cell growth and replication, and topoisomerases, which break and rejoin nucleic acid strands during topological rearrangement of DNA. A Tyr-DNA phosphodiesterase functions in DNA repair by hydrolyzing dead-end covalent intermediates formed between

35 topoisomerase I and DNA (Pouliot, J.J. et al. (1999) Science 286:552-555; Yang, S.-W. (1996) Proc.

Natl. Acad. Sci. USA 93:11534-11539).

Acid sphingomyelinase is a phosphodiesterase which hydrolyzes the membrane phospholipid sphingomyelin to produce ceramide and phosphorylcholine. Phosphorylcholine is used in the synthesis of phosphatidylcholine, which is involved in numerous intracellular signaling pathways, while ceramide is an essential precursor for the generation of gangliosides, membrane lipids found in high concentration in neural tissue. Defective acid sphingomyelinase leads to a build-up of sphingomyelin molecules in lysosomes, resulting in Niemann-Pick disease (Schuchman, E.H. and S.R. Miranda (1997) Genet. Test. 1:13-19).

Glycerophosphoryl diester phosphodiesterase (also known as glycerophosphodiester phosphodiesterase) is a phosphodiesterase which hydrolyzes deacetylated phospholipid glycerophosphodiester to produce sn-glycerol-3-phosphate and an alcohol. Glycerophosphocholine, glycerophosphoethanolamine, glycerophosphoglycerol, and glycerophosphoinositol are examples of substrates for glycerophosphoryl diester phosphodiesterases. A glycerophosphoryl diester phosphodiesterase from *E. coli* has broad specificity for glycerophosphodiester substrates (Larson, T.J. et al. (1983) J. Biol. Chem. 248:5428-5432).

Cyclic nucleotide phosphodiesterases (PDEs) are crucial enzymes in the regulation of the cyclic nucleotides cAMP and cGMP. cAMP and cGMP function as intracellular second messengers to transduce a variety of extracellular signals including hormones, light, and neurotransmitters. PDEs degrade cyclic nucleotides to their corresponding monophosphates, thereby regulating the intracellular concentrations of cyclic nucleotides and their effects on signal transduction. Due to their roles as regulators of signal transduction, PDEs have been extensively studied as chemotherapeutic targets (Perry, M.J. and G.A. Higgs (1998) Curr. Opin. Chem. Biol. 2:472-481; Torphy, J.T. (1998) Am. J. Resp. Crit. Care Med. 157:351-370).

Families of mammalian PDEs have been classified based on their substrate specificity and affinity, sensitivity to cofactors, and sensitivity to inhibitory agents (Beavo, J.A. (1995) Physiol. Rev. 75:725-748; Conti, M. et al. (1995) Endocrine Rev. 16:370-389). Several of these families contain distinct genes, many of which are expressed in different tissues as splice variants. Within PDE families, there are multiple isozymes and multiple splice variants of these isozymes (Conti, M. and S.-L.C. Jin (1999) Prog. Nucleic Acid Res. Mol. Biol. 63:1-38). The existence of multiple PDE families, isozymes, and splice variants is an indication of the variety and complexity of the regulatory pathways involving cyclic nucleotides (Houslay, M.D. and G. Milligan (1997) Trends Biochem. Sci. 22:217-224).

Type 1 PDEs (PDE1s) are  $\text{Ca}^{2+}$ /calmodulin-dependent and appear to be encoded by at least three different genes, each having at least two different splice variants (Kakkar, R. et al. (1999) Cell Mol. Life Sci. 55:1164-1186). PDE1s have been found in the lung, heart, and brain. Some PDE1

isozymes are regulated in vitro by phosphorylation/dephosphorylation. Phosphorylation of these PDE1 isozymes decreases the affinity of the enzyme for calmodulin, decreases PDE activity, and increases steady state levels of cAMP (Kakkar, supra). PDE1s may provide useful therapeutic targets for disorders of the central nervous system, and the cardiovascular and immune systems due to the involvement of PDE1s in both cyclic nucleotide and calcium signaling (Perry, M.J. and G.A. Higgs (1998) *Curr. Opin. Chem. Biol.* 2:472-481).

PDE2s are cGMP-stimulated PDEs that have been found in the cerebellum, neocortex, heart, kidney, lung, pulmonary artery, and skeletal muscle (Sadhu, K. et al. (1999) *J. Histochem. Cytochem.* 47:895-906). PDE2s are thought to mediate the effects of cAMP on catecholamine secretion, participate in the regulation of aldosterone (Beavo, supra), and play a role in olfactory signal transduction (Juifls, D.M. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:3388-3395).

PDE3s have high affinity for both cGMP and cAMP, and so these cyclic nucleotides act as competitive substrates for PDE3s. PDE3s play roles in stimulating myocardial contractility, inhibiting platelet aggregation, relaxing vascular and airway smooth muscle, inhibiting proliferation of T-lymphocytes and cultured vascular smooth muscle cells, and regulating catecholamine-induced release of free fatty acids from adipose tissue. The PDE3 family of phosphodiesterases are sensitive to specific inhibitors such as cilostamide, enoximone, and lixazinone. Isozymes of PDE3 can be regulated by cAMP-dependent protein kinase, or by insulin-dependent kinases (Degerman, E. et al. (1997) *J. Biol. Chem.* 272:6823-6826).

PDE4s are specific for cAMP; are localized to airway smooth muscle, the vascular endothelium, and all inflammatory cells; and can be activated by cAMP-dependent phosphorylation. Since elevation of cAMP levels can lead to suppression of inflammatory cell activation and to relaxation of bronchial smooth muscle, PDE4s have been studied extensively as possible targets for novel anti-inflammatory agents, with special emphasis placed on the discovery of asthma treatments. PDE4 inhibitors are currently undergoing clinical trials as treatments for asthma, chronic obstructive pulmonary disease, and atopic eczema. All four known isozymes of PDE4 are susceptible to the inhibitor rolipram, a compound which has been shown to improve behavioral memory in mice (Barad, M. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:15020-15025). PDE4 inhibitors have also been studied as possible therapeutic agents against acute lung injury, endotoxemia, rheumatoid arthritis, multiple sclerosis, and various neurological and gastrointestinal indications (Doherty, A.M. (1999) *Curr. Opin. Chem. Biol.* 3:466-473).

PDE5 is highly selective for cGMP as a substrate (Turko, I.V. et al. (1998) *Biochemistry* 37:4200-4205), and has two allosteric cGMP-specific binding sites (McAllister-Lucas, L.M. et al. (1995) *J. Biol. Chem.* 270:30671-30679). Binding of cGMP to these allosteric binding sites seems to be important for phosphorylation of PDE5 by cGMP-dependent protein kinase rather than for direct

regulation of catalytic activity. High levels of PDE5 are found in vascular smooth muscle, platelets, lung, and kidney. The inhibitor zaprinast is effective against PDE5 and PDE1s. Modification of zaprinast to provide specificity against PDE5 has resulted in sildenafil (VIAGRA; Pfizer, Inc., New York NY), a treatment for male erectile dysfunction (Terrett, N. et al. (1996) *Bioorg. Med. Chem.*

5 Lett. 6:1819-1824). Inhibitors of PDE5 are currently being studied as agents for cardiovascular therapy (Perry, M.J. and G.A. Higgs (1998) *Curr. Opin. Chem. Biol.* 2:472-481).

PDE6s, the photoreceptor cyclic nucleotide phosphodiesterases, are crucial components of the phototransduction cascade. In association with the G-protein transducin, PDE6s hydrolyze cGMP to regulate cGMP-gated cation channels in photoreceptor membranes. In addition to the cGMP-  
10 binding active site, PDE6s also have two high-affinity cGMP-binding sites which are thought to play a regulatory role in PDE6 function (Artemyev, N.O. et al. (1998) *Methods* 14:93-104). Defects in PDE6s have been associated with retinal disease. Retinal degeneration in the rd mouse (Yan, W. et al. (1998) *Invest. Ophthalmol. Vis. Sci.* 39:2529-2536), autosomal recessive retinitis pigmentosa in humans (Danciger, M. et al. (1995) *Genomics* 30:1-7), and rod/cone dysplasia 1 in Irish Setter dogs  
15 (Suber, M.L. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:3968-3972) have been attributed to mutations in the PDE6B gene.

The PDE7 family of PDEs consists of only one known member having multiple splice variants (Bloom, T.J. and J.A. Beavo (1996) *Proc. Natl. Acad. Sci. USA* 93:14188-14192). PDE7s are cAMP specific, but little else is known about their physiological function. Although mRNAs  
20 encoding PDE7s are found in skeletal muscle, heart, brain, lung, kidney, and pancreas, expression of PDE7 proteins is restricted to specific tissue types (Han, P. et al. (1997) *J. Biol. Chem.* 272:16152-16157; Perry, M.J. and G.A. Higgs (1998) *Curr. Opin. Chem. Biol.* 2:472-481). PDE7s are very closely related to the PDE4 family; however, PDE7s are not inhibited by rolipram, a specific inhibitor of PDE4s (Beavo, supra).

25 PDE8s are cAMP specific, and are closely related to the PDE4 family. PDE8s are expressed in thyroid gland, testis, eye, liver, skeletal muscle, heart, kidney, ovary, and brain. The cAMP-hydrolyzing activity of PDE8s is not inhibited by the PDE inhibitors rolipram, vinpocetine, milrinone, IBMX (3-isobutyl-1-methylxanthine), or zaprinast, but PDE8s are inhibited by dipyridamole (Fisher, D.A. et al. (1998) *Biochem. Biophys. Res. Commun.* 246:570-577; Hayashi, M. et al. (1998)  
30 *Biochem. Biophys. Res. Commun.* 250:751-756; Soderling, S.H. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:8991-8996).

PDE9s are cGMP specific and most closely resemble the PDE8 family of PDEs. PDE9s are expressed in kidney, liver, lung, brain, spleen, and small intestine. PDE9s are not inhibited by sildenafil (VIAGRA; Pfizer, Inc., New York NY), rolipram, vinpocetine, dipyridamole, or IBMX (3-  
35 isobutyl-1-methylxanthine), but they are sensitive to the PDE5 inhibitor zaprinast (Fisher, D.A. et al.



(1998) J. Biol. Chem. 273:15559-15564; Soderling, S.H. et al. (1998) J. Biol. Chem. 273:15553-15558).

PDE10s are dual-substrate PDEs, hydrolyzing both cAMP and cGMP. PDE10s are expressed in brain, thyroid, and testis. (Soderling, S.H. et al. (1999) Proc. Natl. Acad. Sci. USA 96:7071-7076; Fujishige, K. et al. (1999) J. Biol. Chem. 274:18438-18445; Loughney, K. et al (1999) Gene 234:109-117).

PDEs are composed of a catalytic domain of about 270-300 amino acids, an N-terminal regulatory domain responsible for binding cofactors, and, in some cases, a hydrophilic C-terminal domain of unknown function (Conti, M. and S.-L.C. Jin (1999) Prog. Nucleic Acid Res. Mol. Biol., 63:1-38). A conserved, putative zinc-binding motif, HDXXHXGXXN, has been identified in the catalytic domain of all PDEs. N-terminal regulatory domains include non-catalytic cGMP-binding domains in PDE2s, PDE5s, and PDE6s; calmodulin-binding domains in PDE1s; and domains containing phosphorylation sites in PDE3s and PDE4s. In PDE5, the N-terminal cGMP-binding domain spans about 380 amino acid residues and comprises tandem repeats of the conserved sequence motif N(R/K)XnFX<sub>3</sub>DE (McAllister-Lucas, L.M. et al. (1993) J. Biol. Chem. 268:22863-22873). The NKXnD motif has been shown by mutagenesis to be important for cGMP binding (Turko, I.V. et al. (1996) J. Biol. Chem. 271:22240-22244). PDE families display approximately 30% amino acid identity within the catalytic domain; however, isozymes within the same family typically display about 85-95% identity in this region (e.g. PDE4A vs PDE4B). Furthermore, within a family there is extensive similarity (>60%) outside the catalytic domain; while across families, there is little or no sequence similarity outside this domain.

Many of the constituent functions of immune and inflammatory responses are inhibited by agents that increase intracellular levels of cAMP (Verghese, M.W. et al. (1995) Mol. Pharmacol. 47:1164-1171). A variety of diseases have been attributed to increased PDE activity and associated with decreased levels of cyclic nucleotides. For example, a form of diabetes insipidus in mice has been associated with increased PDE4 activity, an increase in low-K<sub>m</sub> cAMP PDE activity has been reported in leukocytes of atopic patients, and PDE3 has been associated with cardiac disease.

Many inhibitors of PDEs have been identified and have undergone clinical evaluation (Perry, M.J. and G.A. Higgs (1998) Curr. Opin. Chem. Biol. 2:472-481; Torphy, T.J. (1998) Am. J. Respir. Crit. Care Med. 157:351-370). PDE3 inhibitors are being developed as antithrombotic agents, antihypertensive agents, and as cardiotonic agents useful in the treatment of congestive heart failure. Rolipram, a PDE4 inhibitor, has been used in the treatment of depression, and other inhibitors of PDE4 are undergoing evaluation as anti-inflammatory agents. Rolipram has also been shown to inhibit lipopolysaccharide (LPS) induced TNF- $\alpha$  which has been shown to enhance HIV-1 replication in vitro. Therefore, rolipram may inhibit HIV-1 replication (Angel, J.B. et al. (1995) AIDS

9:1137-1144). Additionally, rolipram, based on its ability to suppress the production of cytokines such as TNF- $\alpha$  and  $\beta$  and interferon  $\gamma$ , has been shown to be effective in the treatment of encephalomyelitis. Rolipram may also be effective in treating tardive dyskinesia and was effective in treating multiple sclerosis in an experimental animal model (Sommer, N. et al. (1995) Nat. Med.

5 1:244-248; Sasaki, H. et al. (1995) Eur. J. Pharmacol. 282:71-76).

Theophylline is a nonspecific PDE inhibitor used in the treatment of bronchial asthma and other respiratory diseases. Theophylline is believed to act on airway smooth muscle function and in an anti-inflammatory or immunomodulatory capacity in the treatment of respiratory diseases (Banner, K.H. and C.P. Page (1995) Eur. Respir. J. 8:996-1000). Pentoxifylline is another nonspecific PDE  
10 inhibitor used in the treatment of intermittent claudication and diabetes-induced peripheral vascular disease. Pentoxifylline is also known to block TNF- $\alpha$  production and may inhibit HIV-1 replication (Angel et al., supra).

PDEs have been reported to affect cellular proliferation of a variety of cell types (Conti et al. (1995) Endocrine Rev. 16:370-389) and have been implicated in various cancers. Growth of prostate  
15 carcinoma cell lines DU145 and LNCaP was inhibited by delivery of cAMP derivatives and PDE inhibitors (Bang, Y.J. et al. (1994) Proc. Natl. Acad. Sci. USA 91:5330-5334). These cells also showed a permanent conversion in phenotype from epithelial to neuronal morphology. It has also been suggested that PDE inhibitors have the potential to regulate mesangial cell proliferation (Matousovich, K. et al. (1995) J. Clin. Invest. 96:401-410) and lymphocyte proliferation (Joulain, C. et  
20 al. (1995) J. Lipid Mediat. Cell Signal. 11:63-79). A cancer treatment has been described that involves intracellular delivery of PDEs to particular cellular compartments of tumors, resulting in cell death (Deonarain, M.P. and A.A. Epenetos (1994) Br. J. Cancer 70:786-794).

#### Phosphotriesterases

Phosphotriesterases (PTE, paraoxonases) are enzymes that hydrolyze toxic organophosphorus  
25 compounds and have been isolated from a variety of tissues. The enzymes appear to be lacking in birds and insects and abundant in mammals, explaining the reduced tolerance of birds and insects to organophosphorus compound (Vilanova, E. and Sogorb, M.A. (1999) Crit. Rev. Toxicol. 29:21-57). Phosphotriesterases play a central role in the detoxification of insecticides by mammals. Phosphotriesterase activity varies among individuals and is lower in infants than adults. Knockout  
30 mice are markedly more sensitive to the organophosphate-based toxins diazoxon and chlorpyrifos oxon (Furlong, C.E., et al. (2000) Neurotoxicology 21:91-100). PTEs have attracted interest as enzymes capable of the detoxification of organophosphate-containing chemical waste and warfare reagents (e.g., parathion), in addition to pesticides and insecticides. Some studies have also implicated phosphotriesterase in atherosclerosis and diseases involving lipoprotein metabolism.

#### Thioesterases

35

Two soluble thioesterases involved in fatty acid biosynthesis have been isolated from mammalian tissues, one which is active only toward long-chain fatty-acyl thioesters and one which is active toward thioesters with a wide range of fatty-acyl chain-lengths. These thioesterases catalyze the chain-terminating step in the *de novo* biosynthesis of fatty acids. Chain termination involves the hydrolysis of the thioester bond which links the fatty acyl chain to the 4'-phosphopantetheine prosthetic group of the acyl carrier protein (ACP) subunit of the fatty acid synthase (Smith, S. (1981a) Methods Enzymol. 71:181-188; Smith, S. (1981b) Methods Enzymol. 71:188-200).

E. coli contains two soluble thioesterases, thioesterase I which is active only toward long-chain acyl thioesters, and thioesterase II (TEII) which has a broad chain-length specificity (Naggert, J. et al. (1991) J. Biol. Chem. 266:11044-11050). E. coli TEII does not exhibit sequence similarity with either of the two types of mammalian thioesterases which function as chain-terminating enzymes in *de novo* fatty acid biosynthesis. Unlike the mammalian thioesterases, E. coli TEII lacks the characteristic serine active site gly-X-ser-X-gly sequence motif and is not inactivated by the serine modifying agent diisopropyl fluorophosphate. However, modification of histidine 58 by iodoacetamide and diethylpyrocarbonate abolished TEII activity. Overexpression of TEII did not alter fatty acid content in E. coli, which suggests that it does not function as a chain-terminating enzyme in fatty acid biosynthesis (Naggert et al., *supra*). For that reason, Naggert et al. (*supra*) proposed that the physiological substrates for E. coli TEII may be coenzyme A (CoA)-fatty acid esters instead of ACP-phosphopantetheine-fatty acid esters.

#### Carboxylesterases

Mammalian carboxylesterases constitute a multigene family expressed in a variety of tissues and cell types. Isozymes have significant sequence homology and are classified primarily on the basis of amino acid sequence. Acetylcholinesterase, butyrylcholinesterase, and carboxylesterase are grouped into the serine super family of esterases (B-esterases). Other carboxylesterases included thyroglobulin, thrombin, Factor IX, gliotactin, and plasminogen. Carboxylesterases catalyze the hydrolysis of ester- and amide- groups from molecules and are involved in detoxification of drugs, environmental toxins, and carcinogens. Substrates for carboxylesterases include short- and long-chain acyl-glycerols, acylcarnitine, carbonates, dipivefrin hydrochloride, cocaine, salicylates, capsaicin, palmitoyl-coenzyme A, imidapril, haloperidol, pyrrolizidine alkaloids, steroids, p-nitrophenyl acetate, malathion, butanilcaine, and isocarboxazide. The enzymes often demonstrate low substrate specificity. Carboxylesterases are also important for the conversion of prodrugs to their respective free acids, which may be the active form of the drug (e.g., lovastatin, used to lower blood cholesterol) (reviewed in Satoh, T. and Hosokawa, M. (1998) Annu. Rev. Pharmacol. Toxicol.38:257-288).

Neuroligins are a class of molecules that (i) have N-terminal signal sequences, (ii) resemble

cell-surface receptors, (iii) contain carboxylesterase domains, (iv) are highly expressed in the brain, and (v) bind to neurexins in a calcium-dependent manner. Despite the homology to carboxylesterases, neuroligins lack the active site serine residue, implying a role in substrate binding rather than catalysis (Ichtchenko, K. et al. (1996) J. Biol. Chem. 271:2676-2682).

#### 5 Squalene epoxidase

Squalene epoxidase (squalene monooxygenase, SE) is a microsomal membrane-bound, FAD-dependent oxidoreductase that catalyzes the first oxygenation step in the sterol biosynthetic pathway of eukaryotic cells. Cholesterol is an essential structural component of cytoplasmic membranes acquired via the LDL receptor-mediated pathway or the biosynthetic pathway. In the latter case, all  
10 27 carbon atoms in the cholesterol molecule are derived from acetyl-CoA (Stryer, L., supra). SE converts squalene to 2,3(*S*)-oxidosqualene, which is then converted to lanosterol and then cholesterol. The steps involved in cholesterol biosynthesis are summarized below (Stryer, L (1988) Biochemistry. W.H Freeman and Co., Inc. New York. pp. 554-560 and Sakakibara, J. et al. (1995) 270:17-20): acetate (from Acetyl-CoA) → 3-hydroxy-3-methyl-glutaryl CoA → mevalonate → 5-phosphomevalonate  
15 → 5-pyrophosphomevalonate → isopentenyl pyrophosphate → dimethylallyl pyrophosphate → geranyl pyrophosphate → farnesyl pyrophosphate → squalene → squalene epoxide → lanosterol → cholesterol

While cholesterol is essential for the viability of eukaryotic cells, inordinately high serum cholesterol levels results in the formation of atherosclerotic plaques in the arteries of higher organisms. This deposition of highly insoluble lipid material onto the walls of essential blood vessels  
20 (e.g., coronary arteries) results in decreased blood flow and potential necrosis of the tissues deprived of adequate blood flow. HMG-CoA reductase is responsible for the conversion of 3-hydroxyl-3-methyl-glutaryl CoA (HMG-CoA) to mevalonate, which represents the first committed step in cholesterol biosynthesis. HMG-CoA is the target of a number of pharmaceutical compounds designed to lower plasma cholesterol levels. However, inhibition of MHG-CoA also results in the  
25 reduced synthesis of non-sterol intermediates (e.g., mevalonate) required for other biochemical pathways. SE catalyzes a rate-limiting reaction that occurs later in the sterol synthesis pathway and cholesterol in the only end product of the pathway following the step catalyzed by SE. As a result, SE is the ideal target for the design of anti-hyperlipidemic drugs that do not cause a reduction in other necessary intermediates (Nakamura, Y. et al. (1996) 271:8053-8056).

#### 30 Epoxide hydrolases

Epoxide hydrolases catalyze the addition of water to epoxide-containing compounds, thereby hydrolyzing epoxides to their corresponding 1,2-diols. They are related to bacterial haloalkane dehalogenases and show sequence similarity to other members of the  $\alpha/\beta$  hydrolase fold family of enzymes (e.g., bromoperoxidase A2 from Streptomyces aureofaciens, hydroxymuconic semialdehyde  
35 hydrolases from Pseudomonas putida, and haloalkane dehalogenase from Xanthobacter

autotrophicus). Epoxide hydrolases are ubiquitous in nature and have been found in mammals, invertebrates, plants, fungi, and bacteria. This family of enzymes is important for the detoxification of xenobiotic epoxide compounds which are often highly electrophilic and destructive when introduced into an organism. Examples of epoxide hydrolase reactions include the hydrolysis of

5 cis-9,10-epoxyoctadec-9(Z)-enoic acid (leukotoxin) to form its corresponding diol, threo-9,10-dihydroxyoctadec-12(Z)-enoic acid (leukotoxin diol), and the hydrolysis of cis-12,13-epoxyoctadec-9(Z)-enoic acid (isoleukotoxin) to form its corresponding diol threo-12,13-dihydroxyoctadec-9(Z)-enoic acid (isoleukotoxin diol). Leukotoxins alter membrane permeability and ion transport and cause inflammatory responses. In addition, epoxide carcinogens

10 are known to be produced by cytochrome P450 as intermediates in the detoxification of drugs and environmental toxins.

The enzymes possess a catalytic triad composed of Asp (the nucleophile), Asp (the histidine-supporting acid), and His (the water-activating histidine). The reaction mechanism of epoxide hydrolase proceeds via a covalently bound ester intermediate initiated by the nucleophilic

15 attack of one of the Asp residues on the primary carbon atom of the epoxide ring of the target molecule, leading to a covalently bound ester intermediate (Michael Arand, M. et al. (1996) J. Biol. Chem. 271:4223-4229; Rink, R. et al. (1997) J. Biol. Chem. 272:14650-14657; Argiriadi, M.A. et al. (2000) J. Biol. Chem. 275:15265-15270).

#### Enzymes involved in tyrosine catalysis

20 The degradation of the amino acid tyrosine to either succinate and pyruvate or fumarate and acetoacetate, requires a large number of enzymes and generates a large number of intermediate compounds. In addition, many xenobiotic compounds may be metabolized using one or more reactions that are part of the tyrosine catabolic pathway. While the pathway has been studied primarily in bacteria, tyrosine degradation is known to occur in a variety of organisms and is likely to

25 involve many of the same biological reactions.

The enzymes involved in the degradation of tyrosine to succinate and pyruvate (e.g., in *Arthrobacter* species) include 4-hydroxyphenylpyruvate oxidase, 4-hydroxyphenylacetate 3-hydroxylase, 3,4-dihydroxyphenylacetate 2,3-dioxygenase, 5-carboxymethyl-2-hydroxymuconic semialdehyde dehydrogenase, *trans,cis*-5-carboxymethyl-2-hydroxymuconate isomerase,

30 homoprotocatechuate isomerase/decarboxylase, *cis*-2-oxohept-3-ene-1,7-dioate hydratase, 2,4-dihydroxyhept-*trans*-2-ene-1,7-dioate aldolase, and succinic semialdehyde dehydrogenase.

The enzymes involved in the degradation of tyrosine to fumarate and acetoacetate (e.g., in *Pseudomonas* species) include 4-hydroxyphenylpyruvate dioxygenase, homogentisate 1,2-dioxygenase, maleylacetoacetate isomerase, and fumarylacetoacetase. 4-hydroxyphenylacetate

35 1-hydroxylase may also be involved if intermediates from the succinate/pyruvate pathway are

accepted.

Additional enzymes associated with tyrosine metabolism in different organisms include 4-chlorophenylacetate-3,4-dioxygenase, aromatic aminotransferase, 5-oxopent-3-ene-1,2,5-tricarboxylate decarboxylase, 2-oxo-hept-3-ene-1,7-dioate hydratase, and 5-carboxymethyl-2-hydroxymuconate isomerase (Ellis, L.B.M. et al. (1999) *Nucleic Acids Res.* 27:373-376; Wackett, L.P. and Ellis, L.B.M. (1996) *J. Microbiol. Meth.* 25:91-93; and Schmidt, M. (1996) *Amer. Soc. Microbiol. News* 62:102).

In humans, acquired or inherited genetic defects in enzymes of the tyrosine degradation pathway may result in hereditary tyrosinemia. One form of this disease, hereditary tyrosinemia I (HT1) is caused by a deficiency in the enzyme fumarylacetoacetate hydrolase, the last enzyme in the pathway in organisms that metabolize tyrosine to fumarate and acetoacetate. HT1 is characterized by progressive liver damage beginning at infancy, and increased risk for liver cancer (Endo, F. et al. (1997) *J. Biol. Chem.* 272:24426-24432).

The discovery of new drug metabolizing enzymes and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of autoimmune/inflammatory, cell proliferative, developmental, endocrine, eye, metabolic, and gastrointestinal disorders, including liver disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of drug metabolizing enzymes.

## SUMMARY OF THE INVENTION

The invention features purified polypeptides, drug metabolizing enzymes, referred to collectively as "DME" and individually as "DME-1," "DME-2," "DME-3," "DME-4," "DME-5," "DME-6," "DME-7," "DME-8," "DME-9," and "DME-10." In one aspect, the invention provides an isolated polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-10.

The invention further provides an isolated polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, b) a naturally occurring polypeptide comprising an amino acid

sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10.

- 5 In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-10. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:11-20.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide selected from the group  
10 consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, and d) an immunogenic fragment of a polypeptide having an  
15 amino acid sequence selected from the group consisting of SEQ ID NO:1-10. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting  
20 of SEQ ID NO:1-10, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10. The method comprises  
25 a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid  
30 sequence selected from the group consisting of SEQ ID NO:1-10, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the  
35 group consisting of SEQ ID NO:1-10.

The invention further provides an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:11-20, b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:11-20, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:11-20, b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:11-20, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:11-20, b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:11-20, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence



selected from the group consisting of SEQ ID NO:1-10, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, and a pharmaceutically acceptable excipient. In one  
5 embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-10. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional DME, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an  
10 agonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, c) a biologically active fragment of a  
15 polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of  
20 treating a disease or condition associated with decreased expression of functional DME, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, b) a naturally occurring  
25 polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10. The method comprises a) exposing a sample comprising the  
30 polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional DME, comprising administering to a patient in need of such treatment the composition.

35 The invention further provides a method of screening for a compound that specifically binds

to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:11-20, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:11-20, ii) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:11-20, iii) a polynucleotide having a sequence complementary to i), iv) a polynucleotide complementary to the

polynucleotide of ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:11-20, ii) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:11-20, iii) a polynucleotide complementary to the polynucleotide of i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

## BRIEF DESCRIPTION OF THE TABLES

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the present invention.

Table 2 shows the GenBank identification number and annotation of the nearest GenBank homolog for polypeptides of the invention. The probability score for the match between each polypeptide and its GenBank homolog is also shown.

Table 3 shows structural features of polypeptide sequences of the invention, including predicted motifs and domains, along with the methods, algorithms, and searchable databases used for analysis of the polypeptides.

Table 4 lists the cDNA and genomic DNA fragments which were used to assemble polynucleotide sequences of the invention, along with selected fragments of the polynucleotide sequences.

Table 5 shows the representative cDNA library for polynucleotides of the invention.

Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

Table 7 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

## DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is

understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

5 It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

10 Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and  
15 disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

#### DEFINITIONS

"DME" refers to the amino acid sequences of substantially purified DME obtained from any  
20 species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of DME. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of DME either by directly interacting with  
25 DME or by acting on components of the biological pathway in which DME participates.

An "allelic variant" is an alternative form of the gene encoding DME. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to  
30 allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding DME include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as DME or a  
35 polypeptide with at least one functional characteristic of DME. Included within this definition are

polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding DME, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding DME. The encoded protein may also be "altered," and may contain deletions, insertions, or  
5 substitutions of amino acid residues which produce a silent change and result in a functionally equivalent DME. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of DME is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged  
10 amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide,  
15 polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

20 "Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of DME. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates,  
25 small molecules, or any other compound or composition which modulates the activity of DME either by directly interacting with DME or by acting on components of the biological pathway in which DME participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding an epitopic  
30 determinant. Antibodies that bind DME polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin,  
35 thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to

immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic DME, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding DME or fragments of DME may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of, the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
15	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
	Gln	Asn, Glu, His
20	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
25	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
30	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by

an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

5 A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

A "fragment" is a unique portion of DME or the polynucleotide encoding DME which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example,  
10 a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule.

For example, a polypeptide fragment may comprise a certain length of contiguous amino acids  
15 selected from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:11-20 comprises a region of unique polynucleotide sequence that  
20 specifically identifies SEQ ID NO:11-20, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:11-20 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:11-20 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:11-20 and the region of SEQ ID NO:11-20 to which the fragment corresponds are routinely  
25 determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-10 is encoded by a fragment of SEQ ID NO:11-20. A fragment of SEQ ID NO:1-10 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-10. For example, a fragment of SEQ ID NO:1-10 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-  
30 10. The precise length of a fragment of SEQ ID NO:1-10 and the region of SEQ ID NO:1-10 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A  
35 "full length" polynucleotide sequence encodes a "full length" polypeptide sequence.



"Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

*Reward for match: 1*

*Penalty for mismatch: -2*

*Open Gap: 5 and Extension Gap: 2 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 11*

*Filter: on*

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

*Open Gap: 11 and Extension Gap: 1 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 3*

*Filter: on*

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

“Human artificial chromosomes” (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term “humanized antibody” refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

“Hybridization” refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the “washing” step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating  $T_m$  and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%.

5 Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly  
10 under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A  
15 hybridization complex may be formed in solution (e.g., C<sub>0</sub>t or R<sub>0</sub>t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide  
20 sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

25 An "immunogenic fragment" is a polypeptide or oligopeptide fragment of DME which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of DME which is useful in any of the antibody production methods disclosed herein or known in the art.

30 The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of DME. For example, modulation  
35 may cause an increase or a decrease in protein activity, binding characteristics, or any other

biological, functional, or immunological properties of DME.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an DME may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of DME.

"Probe" refers to nucleic acid sequences encoding DME, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

"Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for

example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs  
5 can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to  
10 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from  
15 megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection  
20 programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both  
25 unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

30 A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have  
35 been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently,

a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing DME, nucleic acids encoding DME, or fragments thereof may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

5       A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign  
10       nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed cells" includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as  
15       well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the  
20       art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention  
25       include bacteria, cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

30       A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least  
35       92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least



99% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass, "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length of one of the polypeptides.

## THE INVENTION

The invention is based on the discovery of new human drug metabolizing enzymes (DME), the polynucleotides encoding DME, and the use of these compositions for the diagnosis, treatment, or prevention of autoimmune/inflammatory, cell proliferative, developmental, endocrine, eye, metabolic, and gastrointestinal disorders, including liver disorders.

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is denoted by both a polypeptide sequence identification number (Polypeptide SEQ ID NO:) and an Incyte polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown.

Table 2 shows sequences with homology to the polypeptides of the invention as identified by BLAST analysis against the GenBank protein (genpept) database. Columns 1 and 2 show the

polypeptide sequence identification number (Polypeptide SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for polypeptides of the invention. Column 3 shows the GenBank identification number (Genbank ID NO:) of the nearest GenBank homolog. Column 4 shows the probability score for the match between each polypeptide and its GenBank homolog. Column 5 shows the annotation of the GenBank homolog along with relevant citations where applicable, all of which are expressly incorporated by reference herein.

Table 3 shows various structural features of the polypeptides of the invention. Columns 1 and 2 show the polypeptide sequence identification number (SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Genetics Computer Group, Madison WI). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs. Column 7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

Together, Tables 2 and 3 summarize the properties of polypeptides of the invention, and these properties establish that the claimed polypeptides are drug metabolizing enzymes. For example, SEQ ID NO:9 is 32% identical to human putative N-acetyltransferase Camello 2 (GenBank ID g6651438) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is  $6.50\text{E-}23$ , which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:9 also contains a signal peptide signature sequence as determined by the SPScan program, and an acetyltransferase domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. The probability value of the HMM comparison to acetyltransferase domain is  $3.2\text{e-}12$ . (See Table 3.) Based on BLAST and HMM analyses, the protein of SEQ ID NO:9 is an N-acetyltransferase which N-acetylates aromatic amines and hydrazine-containing compounds including but not limited to para-aminobenzoic acid para-

database (See Table 7 for descriptions). All of these signature sequence identifications are highly statistically significant (see Table 7 for threshold values); for example, the probability value of the HMM comparison to the aldo-keto reductase family signature sequence is  $5.6e-170$ . Based on BLAST, HMM, BLIMPS, MOTIFS, and ProfileScan analyses, the protein of SEQ ID NO:10 is an aldo/keto reductase which reduces carbonyl-containing sugars and aromatic compounds, including, but not limited to, glucose, and carbonyl-containing drug molecules and xenobiotics. SEQ ID NO:1-8 were analyzed and annotated in a similar manner. The algorithms and parameters for the analysis of SEQ ID NO:1-10 are described in Table 7.

As shown in Table 4, the full length polynucleotide sequences of the present invention were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Columns 1 and 2 list the polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and the corresponding Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) for each polynucleotide of the invention. Column 3 shows the length of each polynucleotide sequence in basepairs. Column 4 lists fragments of the polynucleotide sequences which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO:11-20 or that distinguish between SEQ ID NO:11-20 and related polynucleotide sequences. Column 5 shows identification numbers corresponding to cDNA sequences, coding sequences (exons) predicted from genomic DNA, and/or sequence assemblages comprised of both cDNA and genomic DNA. These sequences were used to assemble the full length polynucleotide sequences of the invention. Columns 6 and 7 of Table 4 show the nucleotide start (5') and stop (3') positions of the cDNA and genomic sequences in column 5 relative to their respective full length sequences.

The identification numbers in Column 5 of Table 4 may refer specifically, for example, to Incyte cDNAs along with their corresponding cDNA libraries. For example, 3700065F6 is the identification number of an Incyte cDNA sequence, and SININOT05 is the cDNA library from which it is derived. Incyte cDNAs for which cDNA libraries are not indicated were derived from pooled cDNA libraries (e.g., 71447910V1). Alternatively, the identification numbers in column 5 may refer to GenBank cDNAs or ESTs (e.g., g5663459) which contributed to the assembly of the full length polynucleotide sequences. Alternatively, the identification numbers in column 5 may refer to coding regions predicted by Genscan analysis of genomic DNA. For example, GNN.g7025744\_000055\_002 is the identification number of a Genscan-predicted coding sequence, with g7025744 being the GenBank identification number of the sequence to which Genscan was applied. The Genscan-predicted coding sequences may have been edited prior to assembly. (See Example IV.) Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon stitching" algorithm. For

example, FL152824\_00001 represents a "stitched" sequence in which 152824 is the identification number of the cluster of sequences to which the algorithm was applied, and 00001 is the number of the prediction generated by the algorithm. (See Example V.) Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon-stretching" algorithm. (See Example V.) In some cases, Incyte cDNA coverage redundant with the sequence coverage shown in column 5 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA identification numbers are not shown.

Table 5 shows the representative cDNA libraries for those full length polynucleotide sequences which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to assemble and confirm the above polynucleotide sequences. The tissues and vectors which were used to construct the cDNA libraries shown in Table 5 are described in Table 6.

The invention also encompasses DME variants. A preferred DME variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the DME amino acid sequence, and which contains at least one functional or structural characteristic of DME.

The invention also encompasses polynucleotides which encode DME. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:11-20, which encodes DME. The polynucleotide sequences of SEQ ID NO:11-20, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding DME. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding DME. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:11-20 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:11-20. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of DME.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding DME, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be

produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring DME, and all such variations are to be considered as  
5 being specifically disclosed.

Although nucleotide sequences which encode DME and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring DME under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding DME or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally  
10 occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding DME and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater  
15 half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode DME and DME derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to  
20 introduce mutations into a sequence encoding DME or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:11-20 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.*  
25 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal  
35 cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA

sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding DME may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the

emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments  
5 which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode DME may be cloned in recombinant DNA molecules that direct expression of DME, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a  
10 functionally equivalent amino acid sequence may be produced and used to express DME.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter DME-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and  
15 synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such  
20 as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Crameri, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of DME, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene  
25 variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random  
30 point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

35 In another embodiment, sequences encoding DME may be synthesized, in whole or in part,

using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser. 7*:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser. 7*:225-232.) Alternatively, DME itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques.

5 (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of DME, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide  
10 or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, supra, pp. 28-53.)

15 In order to express a biologically active DME, the nucleotide sequences encoding DME or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in  
20 polynucleotide sequences encoding DME. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding DME. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding DME and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional  
25 or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See,  
30 e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding DME and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989)  
35 Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and



16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding DME. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509; Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945; Takamatsu, N. (1987) EMBO J. 6:307-311; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659; and Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) Cancer Gen. Ther. 5(6):350-356; Yu, M. et al. (1993) Proc. Natl. Acad. Sci. USA 90(13):6340-6344; Buller, R.M. et al. (1985) Nature 317(6040):813-815; McGregor, D.P. et al. (1994) Mol. Immunol. 31(3):219-226; and Verma, I.M. and N. Somia (1997) Nature 389:239-242.) The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding DME. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding DME can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding DME into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of DME are needed, e.g. for the production of antibodies, vectors which direct high level expression of DME may be used. For example, vectors containing the strong, inducible SP6 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of DME. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such

vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; and Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184.)

5 Plant systems may also be used for expression of DME. Transcription of sequences encoding DME may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. 10 (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases 15 where an adenovirus is used as an expression vector, sequences encoding DME may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses DME in host cells. (See, e.g., Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression. 20

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino 25 polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of DME in cell lines is preferred. For example, sequences encoding DME can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous 30 expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be 35 propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *apr* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech),  $\beta$  glucuronidase and its substrate  $\beta$ -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding DME is inserted within a marker gene sequence, transformed cells containing sequences encoding DME can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding DME under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding DME and that express DME may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of DME using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on DME is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul

MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding DME include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding DME, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding DME may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode DME may be designed to contain signal sequences which direct secretion of DME through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding DME may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric DME protein containing a heterologous moiety that can be recognized by a commercially available antibody may

facilitate the screening of peptide libraries for inhibitors of DME activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the DME encoding sequence and the heterologous protein sequence, so that DME may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled DME may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, <sup>35</sup>S-methionine.

DME of the present invention or fragments thereof may be used to screen for compounds that specifically bind to DME. At least one and up to a plurality of test compounds may be screened for specific binding to DME. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of DME, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which DME binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express DME, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing DME or cell membrane fractions which contain DME are then contacted with a test compound and binding, stimulation, or inhibition of activity of either DME or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is

detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with DME, either in solution or affixed to a solid support, and detecting the binding of DME to the compound.

Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

DME of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of DME. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for DME activity, wherein DME is combined with at least one test compound, and the activity of DME in the presence of a test compound is compared with the activity of DME in the absence of the test compound. A change in the activity of DME in the presence of the test compound is indicative of a compound that modulates the activity of DME. Alternatively, a test compound is combined with an in vitro or cell-free system comprising DME under conditions suitable for DME activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of DME may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding DME or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent Number 5,175,383 and U.S. Patent Number 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding DME may also be manipulated in vitro in ES cells derived from

human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding DME can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding DME is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress DME, e.g., by secreting DME in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

### THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of DME and drug metabolizing enzymes. In addition, the expression of DME is closely associated with a liver tumor cell line, liver tumor tissue, pancreatic tissue, pituitary gland tissue, brain tissue, small intestine tissue, fetal brain tissue, and allocortex brain tissue. Therefore, DME appears to play a role in autoimmune/inflammatory, cell proliferative, developmental, endocrine, eye, metabolic, and gastrointestinal disorders, including liver disorders. In the treatment of disorders associated with increased DME expression or activity, it is desirable to decrease the expression or activity of DME. In the treatment of disorders associated with decreased DME expression or activity, it is desirable to increase the expression or activity of DME.

Therefore, in one embodiment, DME or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of DME. Examples of such disorders include, but are not limited to, an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome,

systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; a developmental disorder, such as renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, and sensorineural hearing loss; an endocrine disorder, such as disorders of the hypothalamus and pituitary resulting from lesions such as primary brain tumors, adenomas, infarction associated with pregnancy, hypophysectomy, aneurysms, vascular malformations, thrombosis, infections, immunological disorders, and complications due to head trauma; disorders associated with hypopituitarism including hypogonadism, Sheehan syndrome, diabetes insipidus, Kallman's disease, Hand-Schuller-Christian disease, Letterer-Siwe disease, sarcoidosis, empty sella syndrome, and dwarfism; disorders associated with hyperpituitarism including acromegaly, gigantism, and syndrome of inappropriate antidiuretic hormone (ADH) secretion (SIADH) often caused by benign adenoma; disorders associated with hypothyroidism including goiter, myxedema, acute thyroiditis associated with bacterial infection, subacute thyroiditis associated with viral infection, autoimmune thyroiditis (Hashimoto's disease), and cretinism; disorders associated with hyperthyroidism including thyrotoxicosis and its various forms, Grave's disease, pretibial myxedema, toxic multinodular goiter, thyroid carcinoma, and Plummer's disease; disorders associated with hyperparathyroidism including Conn disease (chronic hypercalcemia); pancreatic disorders such as Type I or Type II diabetes mellitus and associated complications; disorders associated with the adrenals such as hyperplasia, carcinoma, or adenoma of the adrenal cortex, hypertension associated with alkalosis, amyloidosis, hypokalemia, Cushing's disease, Liddle's syndrome, and Arnold-Healy-Gordon syndrome, pheochromocytoma tumors, and Addison's disease; disorders associated with gonadal steroid



hormones such as: in women, abnormal prolactin production, infertility, endometriosis, perturbations of the menstrual cycle, polycystic ovarian disease, hyperprolactinemia, isolated gonadotropin deficiency, amenorrhea, galactorrhea, hermaphroditism, hirsutism and virilization, breast cancer, and, in post-menopausal women, osteoporosis; and, in men, Leydig cell deficiency, male climacteric phase, and germinal cell aplasia, hypergonadal disorders associated with Leydig cell tumors, androgen resistance associated with absence of androgen receptors, syndrome of 5  $\alpha$ -reductase, and gynecomastia; an eye disorder, such as conjunctivitis, keratoconjunctivitis sicca, keratitis, episcleritis, iritis, posterior uveitis, glaucoma, amaurosis fugax, ischemic optic neuropathy, optic neuritis, Leber's hereditary optic neuropathy, toxic optic neuropathy, vitreous detachment, retinal detachment, cataract, macular degeneration, central serous chorioretinopathy, retinitis pigmentosa, melanoma of the choroid, retrobulbar tumor, and chiasmal tumor; a metabolic disorder, such as Addison's disease, cerebrotendinous xanthomatosis, congenital adrenal hyperplasia, coumarin resistance, cystic fibrosis, diabetes, fatty hepatocirrhosis, fructose-1,6-diphosphatase deficiency, galactosemia, goiter, glucagonoma, glycogen storage diseases, hereditary fructose intolerance, hyperadrenalism, hypoadrenalism, hyperparathyroidism, hypoparathyroidism, hypercholesterolemia, hyperthyroidism, hypoglycemia, hypothyroidism, hyperlipidemia, hyperlipemia, lipid myopathies, lipodystrophies, lysosomal storage diseases, Menkes syndrome, occipital horn syndrome, mannosidosis, neuraminidase deficiency, obesity, pentosuria phenylketonuria, pseudovitamin D-deficiency rickets; hypocalcemia, hypophosphatemia, and postpubescent cerebellar ataxia, tyrosinemia, and a gastrointestinal disorder, such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, hereditary hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha<sub>1</sub>-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas.

In another embodiment, a vector capable of expressing DME or a fragment or derivative

thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of DME including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified DME in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of DME including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of DME may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of DME including, but not limited to, those listed above.

In a further embodiment, an antagonist of DME may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of DME. Examples of such disorders include, but are not limited to, those autoimmune/inflammatory, cell proliferative, developmental, endocrine, eye, metabolic, and gastrointestinal disorders, including liver disorders, described above. In one aspect, an antibody which specifically binds DME may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express DME.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding DME may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of DME including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of DME may be produced using methods which are generally known in the art. In particular, purified DME may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind DME. Antibodies to DME may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with DME or with any fragment or oligopeptide thereof

which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants  
5 used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to DME have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or  
10 fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of DME amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to DME may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not  
15 limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the  
20 splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce DME-specific  
25 single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as  
30 disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for DME may also be generated. For example, such fragments include, but are not limited to, F(ab')<sub>2</sub> fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of  
35 the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and

easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between DME and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering DME epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for DME. Affinity is expressed as an association constant,  $K_a$ , which is defined as the molar concentration of DME-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The  $K_a$  determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple DME epitopes, represents the average affinity, or avidity, of the antibodies for DME. The  $K_a$  determined for a preparation of monoclonal antibodies, which are monospecific for a particular DME epitope, represents a true measure of affinity. High-affinity antibody preparations with  $K_a$  ranging from about  $10^9$  to  $10^{12}$  L/mole are preferred for use in immunoassays in which the DME-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with  $K_a$  ranging from about  $10^6$  to  $10^7$  L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of DME, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of DME-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding DME, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene

encoding DME. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding DME. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

- 5 In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, 15 R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

- In another embodiment of the invention, polynucleotides encoding DME may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, 25 familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) 30 (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA.* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in DME expression or regulation causes disease, the expression of DME from an appropriate population of transduced cells may alleviate the 35 clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in DME are treated by constructing mammalian expression vectors encoding DME and introducing these vectors by mechanical means into DME-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J-L. and H. Récipon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of DME include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). DME may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or  $\beta$ -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and Blau, H.M. *supra*), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding DME from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to DME expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding DME under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc.*

Natl. Acad. Sci. USA 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) J. Virol. 61:1647-1650; Bender, M.A. et al. (1987) J. Virol. 61:1639-1646; Adam, M.A. and A.D. Miller (1988) J. Virol. 62:3802-3806; Dull, T. et al. (1998) J. Virol. 72:8463-8471; Zufferey, R. et al. (1998) J. Virol. 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4<sup>+</sup> T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) J. Virol. 71:7020-7029; Bauer, G. et al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716; Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding DME to cells which have one or more genetic abnormalities with respect to the expression of DME. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) Annu. Rev. Nutr. 19:511-544 and Verma, I.M. and N. Somia (1997) Nature 389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding DME to target cells which have one or more genetic abnormalities with respect to the expression of DME. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing DME to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 69:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a

cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of  
5 cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to  
10 deliver polynucleotides encoding DME to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA,  
15 resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for DME into the alphavirus genome in place of the capsid-coding region results in the production of a large number of DME-coding RNAs and the synthesis of high levels of DME in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a  
20 persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of DME into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of  
25 manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly,  
30 inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-  
35 177.) A complementary sequence or antisense molecule may also be designed to block translation of



mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example,  
5 engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding DME.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides,  
10 corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared  
15 by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding DME. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively,  
20 these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase  
25 linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

30 An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding DME. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular  
35 chemical entities which are capable of interacting with specific polynucleotide sequences. Effective

compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased DME expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding DME may be therapeutically useful, and in the treatment of disorders associated with  
5 decreased DME expression or activity, a compound which specifically promotes expression of the polynucleotide encoding DME may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective  
10 in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding DME is exposed to at least one test compound thus obtained. The sample  
15 may comprise, for example, an intact or permeabilized cell, or an in vitro cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding DME are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding DME. The amount of hybridization may be quantified, thus forming  
20 the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a Schizosaccharomyces pombe gene  
25 expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a  
30 specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient.

35 Delivery by transfection, by liposome injections, or by polycationic amino polymers may be

achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of DME, antibodies to DME, and mimetics, agonists, antagonists, or inhibitors of DME.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising DME or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, DME or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell

culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

5           A therapeutically effective dose refers to that amount of active ingredient, for example DME or fragments thereof, antibodies of DME, and agonists, antagonists or inhibitors of DME, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the  $ED_{50}$  (the dose therapeutically effective in 50% of the population) or  $LD_{50}$  (the dose  
10   lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the  $LD_{50}/ED_{50}$  ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the  $ED_{50}$  with little or no  
15   toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include  
20   the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1  $\mu\text{g}$  to 100,000  $\mu\text{g}$ , up to a total dose of  
25   about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

### 30   **DIAGNOSTICS**

In another embodiment, antibodies which specifically bind DME may be used for the diagnosis of disorders characterized by expression of DME, or in assays to monitor patients being treated with DME or agonists, antagonists, or inhibitors of DME. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic  
35   assays for DME include methods which utilize the antibody and a label to detect DME in human

body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

5 A variety of protocols for measuring DME, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of DME expression. Normal or standard values for DME expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibodies to DME under conditions suitable for complex formation. The amount of standard complex formation may be  
10 quantitated by various methods, such as photometric means. Quantities of DME expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding DME may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences,  
15 complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of DME may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of DME, and to monitor regulation of DME levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide  
20 sequences, including genomic sequences, encoding DME or closely related molecules may be used to identify nucleic acid sequences which encode DME. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding DME, allelic variants, or related  
25 sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the DME encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:11-20 or from genomic sequences including promoters, enhancers, and introns of the DME gene.

30 Means for producing specific hybridization probes for DNAs encoding DME include the cloning of polynucleotide sequences encoding DME or DME derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a  
35 variety of reporter groups, for example, by radionuclides such as  $^{32}\text{P}$  or  $^{35}\text{S}$ , or by enzymatic labels,

such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding DME may be used for the diagnosis of disorders associated with expression of DME. Examples of such disorders include, but are not limited to, an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS),

- 5 Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis,
- 10 erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura,
- 15 ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers
- 20 including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; a developmental disorder, such as renal tubular acidosis, anemia, Cushing's syndrome,
- 25 achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea
- 30 and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, and sensorineural hearing loss; an endocrine disorder, such as disorders of the hypothalamus and pituitary resulting from lesions such as primary brain tumors, adenomas, infarction associated with pregnancy, hypophysectomy, aneurysms, vascular malformations, thrombosis, infections, immunological disorders, and complications due to head trauma; disorders associated with
- 35 hypopituitarism including hypogonadism, Sheehan syndrome, diabetes insipidus, Kallman's disease,

Hand-Schuller-Christian disease, Letterer-Siwe disease, sarcoidosis, empty sella syndrome, and dwarfism; disorders associated with hyperpituitarism including acromegaly, giantism, and syndrome of inappropriate antidiuretic hormone (ADH) secretion (SIADH) often caused by benign adenoma; disorders associated with hypothyroidism including goiter, myxedema, acute thyroiditis associated with bacterial infection, subacute thyroiditis associated with viral infection, autoimmune thyroiditis (Hashimoto's disease), and cretinism; disorders associated with hyperthyroidism including thyrotoxicosis and its various forms, Grave's disease, pretibial myxedema, toxic multinodular goiter, thyroid carcinoma, and Plummer's disease; disorders associated with hyperparathyroidism including Conn disease (chronic hypercalcemia); pancreatic disorders such as Type I or Type II diabetes mellitus and associated complications; disorders associated with the adrenals such as hyperplasia, carcinoma, or adenoma of the adrenal cortex, hypertension associated with alkalosis, amyloidosis, hypokalemia, Cushing's disease, Liddle's syndrome, and Arnold-Healy-Gordon syndrome, pheochromocytoma tumors, and Addison's disease; disorders associated with gonadal steroid hormones such as: in women, abnormal prolactin production, infertility, endometriosis, perturbations of the menstrual cycle, polycystic ovarian disease, hyperprolactinemia, isolated gonadotropin deficiency, amenorrhea, galactorrhea, hermaphroditism, hirsutism and virilization, breast cancer, and, in post-menopausal women, osteoporosis; and, in men, Leydig cell deficiency, male climacteric phase, and germinal cell aplasia, hypergonadal disorders associated with Leydig cell tumors, androgen resistance associated with absence of androgen receptors, syndrome of 5  $\alpha$ -reductase, and gynecomastia; an eye disorder, such as conjunctivitis, keratoconjunctivitis sicca, keratitis, episcleritis, iritis, posterior uveitis, glaucoma, amaurosis fugax, ischemic optic neuropathy, optic neuritis, Leber's hereditary optic neuropathy, toxic optic neuropathy, vitreous detachment, retinal detachment, cataract, macular degeneration, central serous chorioretinopathy, retinitis pigmentosa, melanoma of the choroid, retrobulbar tumor, and chiasmal tumor; a metabolic disorder, such as Addison's disease, cerebrotendinous xanthomatosis, congenital adrenal hyperplasia, coumarin resistance, cystic fibrosis, diabetes, fatty hepatocirrhosis, fructose-1,6-diphosphatase deficiency, galactosemia, goiter, glucagonoma, glycogen storage diseases, hereditary fructose intolerance, hyperadrenalism, hypoadrenalism, hyperparathyroidism, hypoparathyroidism, hypercholesterolemia, hyperthyroidism, hypoglycemia, hypothyroidism, hyperlipidemia, hyperlipemia, lipid myopathies, lipodystrophies, lysosomal storage diseases, Menkes syndrome, occipital horn syndrome, mannosidosis, neuraminidase deficiency, obesity, pentosuria, phenylketonuria, pseudovitamin D-deficiency rickets; hypocalcemia, hypophosphatemia, and postpubescent cerebellar ataxia, tyrosinemia, and a gastrointestinal disorder, such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric

edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, hereditary hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha<sub>1</sub>-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas. The polynucleotide sequences encoding DME may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered DME expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding DME may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding DME may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding DME in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of DME, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding DME, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.



Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from  
5 several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals  
10 to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding DME may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a  
15 polynucleotide encoding DME, or a fragment of a polynucleotide complementary to the polynucleotide encoding DME, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences  
20 encoding DME may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding DME are used to  
25 amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-  
30 throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA  
35 sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass

spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of DME include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, DME, fragments of DME, or antibodies specific for DME may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance

under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, supra). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for DME to quantify the levels of DME expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendoz, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of

each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding DME may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, for example, Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of

genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding DME on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

5        In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene  
10        discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide  
15        sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

          In another embodiment of the invention, DME, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes  
20        between DME and the agent being tested may be measured.

          Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with DME, or fragments thereof,  
25        and washed. Bound DME is then detected by methods well known in the art. Purified DME can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

          In another embodiment, one may use competitive drug screening assays in which  
30        neutralizing antibodies capable of binding DME specifically compete with a test compound for binding DME. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with DME.

          In additional embodiments, the nucleotide sequences which encode DME may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on  
35        properties of nucleotide sequences that are currently known, including, but not limited to, such

properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure  
5 in any way whatsoever.

The disclosures of all patents, applications and publications, mentioned above and below, including U.S. Ser. No. 60/197,590, U.S. Ser. No. 60/198,403, U.S. Ser. No. 60/200,185, U.S. Ser. No. 60/202,234, and U.S. Ser. No. 60/203,509, are expressly incorporated by reference herein.

10

## EXAMPLES

### I. Construction of cDNA Libraries

15

Incyte cDNAs were derived from cDNA libraries described in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA) and shown in Table 4, column 5. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

20

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)+ RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

25

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g.,  
35 PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), PCDNA2.1 plasmid

(Invitrogen, Carlsbad CA), PBK-CMV plasmid (Stratagene), or pINCY (Incyte Genomics, Palo Alto CA), or derivatives thereof. Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 $\alpha$ , DH10B, or ElectroMAX DH10B from Life Technologies.

## 5 II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8  
10 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal  
15 cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

## 20 III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared  
25 using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI  
30 protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VIII.

The polynucleotide sequences derived from Incyte cDNAs were validated by removing  
35 vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and



programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA sequences or translations thereof were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and hidden Markov model (HMM)-based protein family databases such as PFAM. (HMM is a probabilistic approach which analyzes consensus primary structures of gene families. See, for example, Eddy, S.R. (1996) *Curr. Opin. Struct. Biol.* 6:361-365.) The queries were performed using programs based on BLAST, FASTA, BLIMPS, and HMMER. The Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or Genscan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblages to full length. Assembly was performed using programs based on Phred, Phrap, and Consed, and cDNA assemblages were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length polypeptide sequences. Alternatively, a polypeptide of the invention may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently analyzed by querying against databases such as the GenBank protein databases (genpept), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and hidden Markov model (HMM)-based protein family databases such as PFAM. Full length polynucleotide sequences are also analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments are generated using default parameters specified by the CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and threshold parameters. The first column of Table 7 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value, the greater the identity between two sequences).

The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:11-20. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization

and amplification technologies are described in Table 4, column 4.

#### IV. Identification and Editing of Coding Sequences from Genomic DNA

Putative drug metabolizing enzymes were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., gbpri and gbhtg). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (See Burge, C. and S. Karlin (1997) J. Mol. Biol. 268:78-94, and Burge, C. and S. Karlin (1998) Curr. Opin. Struct. Biol. 8:346-354). The program concatenates predicted exons to form an assembled cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 kb. To determine which of these Genscan predicted cDNA sequences encode drug metabolizing enzymes, the encoded polypeptides were analyzed by querying against PFAM models for drug metabolizing enzymes. Potential drug metabolizing enzymes were also identified by homology to Incyte cDNA sequences that had been annotated as drug metabolizing enzymes. These selected Genscan-predicted sequences were then compared by BLAST analysis to the genpept and gbpri public databases. Where necessary, the Genscan-predicted sequences were then edited by comparison to the top BLAST hit from genpept to correct errors in the sequence predicted by Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing evidence for transcription. When Incyte cDNA coverage was available, this information was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived entirely from edited or unedited Genscan-predicted coding sequences.

#### V. Assembly of Genomic Sequence Data with cDNA Sequence Data

##### "Stitched" Sequences

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification program described in Example IV. Partial cDNAs assembled as described in Example III were mapped to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm based on graph theory and dynamic programming to integrate cDNA and genomic information, generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence. Sequence intervals in which the entire length of the interval was present on more than one sequence in the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present

on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to be brought together, bridged by cDNA sequence. Intervals thus identified were then "stitched" together by the stitching algorithm in the order that they appear along their parent sequences to generate the longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbpri public databases. Incorrect exons predicted by Genscan were corrected by comparison to the top BLAST hit from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

#### **"Stretched" Sequences**

Partial DNA sequences were extended to full length with an algorithm based on BLAST analysis. First, partial cDNAs assembled as described in Example III were queried against public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or GenScan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the translated sequences onto the GenBank protein homolog. Insertions or deletions may occur in the chimeric protein with respect to the original GenBank protein homolog. The GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore "stretched" or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to determine whether it contained a complete gene.

#### **VI. Chromosomal Mapping of DME Encoding Polynucleotides**

The sequences which were used to assemble SEQ ID NO:11-20 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:11-20 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO:, to that map location.

Map locations are represented by ranges, or intervals, of human chromosomes. The map

position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

## VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel (1995) *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum} \{ \text{length}(\text{Seq. 1}), \text{length}(\text{Seq. 2}) \}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end,

or 79% identity and 100% overlap.

Alternatively, polynucleotide sequences encoding DME are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled, at least in part, with overlapping Incyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive system; embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following disease/condition categories: cancer, cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and disease-specific expression of cDNA encoding DME. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA).

#### **VIII. Extension of DME Encoding Polynucleotides**

Full length polynucleotide sequences were also produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was synthesized to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing  $Mg^{2+}$ ,  $(NH_4)_2SO_4$ , and 2-mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage

at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 µl PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 µl of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 µl to 10 µl aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

In like manner, full length polynucleotide sequences are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for such extension, and an appropriate genomic library.

#### **IX. Labeling and Use of Individual Hybridization Probes**

Hybridization probes derived from SEQ ID NO:11-20 are employed to screen cDNAs,

genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250  $\mu$ Ci of [5  $\gamma$ -<sup>32</sup>P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10<sup>7</sup> counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: 10 Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. 15 Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

#### X. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra), 20 mechanical microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Skena (1999), supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding 25 procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Skena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof 30 may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, 35 and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser

desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

## 5 **Tissue or Cell Sample Preparation**

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)<sup>+</sup> RNA is purified using the oligo-(dT) cellulose method. Each poly(A)<sup>+</sup> RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/ $\mu$ l oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/ $\mu$ l RNase inhibitor, 500  $\mu$ M dATP, 500  $\mu$ M dGTP, 500  $\mu$ M dTTP, 40  
10  $\mu$ M dCTP, 40  $\mu$ M dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)<sup>+</sup> RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)<sup>+</sup> RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37°C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium  
15 hydroxide and incubated for 20 minutes at 85°C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments  
20 Inc., Holbrook NY) and resuspended in 14  $\mu$ l 5X SSC/0.2% SDS.

## **Microarray Preparation**

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are  
25 amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5  $\mu$ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water  
30 washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US



Patent No. 5,807,522, incorporated herein by reference. 1  $\mu$ l of the array element DNA, at an average concentration of 100 ng/ $\mu$ l, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene).

- 5 Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60°C followed by washes in 0.2% SDS and distilled water as before.

#### **Hybridization**

- 10 Hybridization reactions contain 9  $\mu$ l of sample mixture consisting of 0.2  $\mu$ g each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm<sup>2</sup> coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the  
15 addition of 140  $\mu$ l of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

#### **Detection**

- 20 Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-  
25 scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores.

- 30 Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

## **XI. Complementary Polynucleotides**

Sequences complementary to the DME-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring DME. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of DME. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the DME-encoding transcript.

## **XII. Expression of DME**

Expression and purification of DME is achieved using bacterial or virus-based expression systems. For expression of DME in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory

element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express DME upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of DME in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding DME by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, DME is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from DME at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified DME obtained by these methods can be used directly in the assays shown in Examples XVI, XVII, and XVIII, where applicable.

### XIII. Functional Assays

DME function is assessed by expressing the sequences encoding DME at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include PCMV SPORT (Life Technologies) and PCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10  $\mu$ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2  $\mu$ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP;

Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of DME on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding DME and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding DME and other genes of interest can be analyzed by northern analysis or microarray techniques.

#### **XIV. Production of DME Specific Antibodies**

DME substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the DME amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using FMOC chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-DME activity by, for example, binding the peptide or DME to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated

goat anti-rabbit IgG.

#### **XV. Purification of Naturally Occurring DME Using Specific Antibodies**

Naturally occurring or recombinant DME is substantially purified by immunoaffinity chromatography using antibodies specific for DME. An immunoaffinity column is constructed by covalently coupling anti-DME antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing DME are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of DME (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/DME binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and DME is collected.

#### **XVI. Identification of Molecules Which Interact with DME**

DME, or biologically active fragments thereof, are labeled with  $^{125}\text{I}$  Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled DME, washed, and any wells with labeled DME complex are assayed. Data obtained using different concentrations of DME are used to calculate values for the number, affinity, and association of DME with the candidate molecules.

Alternatively, molecules interacting with DME are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989) *Nature* 340:245-246, or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

DME may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

#### **XVII. Demonstration of DME Activity**

Cytochrome P450 activity of DME is measured using the 4-hydroxylation of aniline. Aniline is converted to 4-aminophenol by the enzyme, and has an absorption maximum at 630 nm (Gibson and Skett, *supra*). This assay is a convenient measure, but underestimates the total hydroxylation, which also occurs at the 2- and 3- positions. Assays are performed at 37°C and contain an aliquot of the enzyme and a suitable amount of aniline (approximately 2 mM) in reaction buffer. For this reaction, the buffer must contain NADPH or an NADPH-generating cofactor system. One formulation for this reaction buffer includes 85 mM Tris pH 7.4, 15 mM  $\text{MgCl}_2$ , 50 mM nicotinamide, 40 mg trisodium isocitrate, and 2 units isocitrate dehydrogenase, with 8 mg  $\text{NADP}^+$

added to a 10 mL reaction buffer stock just prior to assay. Reactions are carried out in an optical cuvette, and the absorbance at 630 nm is measured. The rate of increase in absorbance is proportional to the enzyme activity in the assay. A standard curve can be constructed using known concentrations of 4-aminophenol.

- 5           1 $\alpha$ ,25-dihydroxyvitamin D 24-hydroxylase activity of DME is determined by monitoring the conversion of <sup>3</sup>H-labeled 1 $\alpha$ ,25-dihydroxyvitamin D (1 $\alpha$ ,25(OH)<sub>2</sub>D) to 24,25-dihydroxyvitamin D (24,25(OH)<sub>2</sub>D) in transgenic rats expressing DME. 1  $\mu$ g of 1 $\alpha$ ,25(OH)<sub>2</sub>D dissolved in ethanol (or ethanol alone as a control) is administered intravenously to approximately 6-week-old male transgenic rats expressing DME or otherwise identical control rats expressing either a defective
- 10           variant of DME or not expressing DME. The rats are killed by decapitation after 8 hrs, and the kidneys are rapidly removed, rinsed, and homogenized in 9 volumes of ice-cold buffer (15 mM Tris-acetate (pH 7.4), 0.19 M sucrose, 2 mM magnesium acetate, and 5 mM sodium succinate). A portion (e.g., 3 ml) of each homogenate is then incubated with 0.25 nM 1 $\alpha$ ,25(OH)<sub>2</sub>[1-<sup>3</sup>H]D, with a specific activity of approximately 3.5 GBq/mmol, for 15 min at 37 °C under oxygen with constant shaking.
- 15           Total lipids are extracted as described (Bligh, E.G. and W.J. Dyer (1959) Can. J. Biochem. Physiol. 37: 911-917) and the chloroform phase is analyzed by HPLC using a FINEPAK SIL column (JASCO, Tokyo, Japan) with a *n*-hexane/chloroform/methanol (10:2.5:1.5) solvent system at a flow rate of 1 ml/min. In the alternative, the chloroform phase is analyzed by reverse phase HPLC using a J SPHERE ODS-AM column (YMC Co. Ltd., Kyoto, Japan) with an acetonitrile buffer system (40
- 20           to 100%, in water, in 30 min) at a flow rate of 1 ml/min. The eluates are collected in fractions of 30 seconds (or less) and the amount of <sup>3</sup>H present in each fraction is measured using a scintillation counter. By comparing the chromatograms of control samples (i.e., samples comprising 1 $\alpha$ ,25-dihydroxyvitamin D or 24,25-dihydroxyvitamin D (24,25(OH)<sub>2</sub>D), with the chromatograms of the reaction products, the relative mobilities of the substrate (1 $\alpha$ ,25(OH)<sub>2</sub>[1-<sup>3</sup>H]D) and product
- 25           (24,25(OH)<sub>2</sub>[1-<sup>3</sup>H]D) are determined and correlated with the fractions collected. The amount of 24,25(OH)<sub>2</sub>[1-<sup>3</sup>H]D produced in control rats is subtracted from that of transgenic rats expressing DME. The difference in the production of 24,25(OH)<sub>2</sub>[1-<sup>3</sup>H]D in the transgenic and control animals is proportional to the amount of 25-hydroxylase activity of DME present in the sample. Confirmation of the identity of the substrate and product(s) is confirmed by means of mass spectroscopy
- 30           (Miyamoto, Y. et al. (1997) J. Biol. Chem. 272:14115-14119).

Flavin-containing monooxygenase activity of DME is measured by chromatographic analysis of metabolic products. For example, Ring, B.J. et al. (1999; Drug Metab. Dis. 27:1099-1103) incubated FMO in 0.1 M sodium phosphate buffer (pH 7.4 or 8.3) and 1 mM NADPH at 37°C, stopped the reaction with an organic solvent, and determined product formation by HPLC.

- 35           Alternatively, activity is measured by monitoring oxygen uptake using a Clark-type electrode. For

example, Ziegler, D.M. and Poulsen, L.L. (1978; Methods Enzymol. 52:142-151) incubated the enzyme at 37°C in an NADPH-generating cofactor system (similar to the one described above) containing the substrate methimazole. The rate of oxygen uptake is proportional to enzyme activity.

UDP glucuronyltransferase activity of DME is measured using a colorimetric determination of free amine groups (Gibson and Skett, supra). An amine-containing substrate, such as 2-aminophenol, is incubated at 37°C with an aliquot of the enzyme in a reaction buffer containing the necessary cofactors (40 mM Tris pH 8.0, 7.5 mM MgCl<sub>2</sub>, 0.025% Triton X-100, 1 mM ascorbic acid, 0.75 mM UDP-glucuronic acid). After sufficient time, the reaction is stopped by addition of ice-cold 20% trichloroacetic acid in 0.1 M phosphate buffer pH 2.7, incubated on ice, and centrifuged to clarify the supernatant. Any unreacted 2-aminophenol is destroyed in this step. Sufficient freshly-prepared sodium nitrite is then added; this step allows formation of the diazonium salt of the glucuronidated product. Excess nitrite is removed by addition of sufficient ammonium sulfamate, and the diazonium salt is reacted with an aromatic amine (for example, N-naphthylethylene diamine) to produce a colored azo compound which can be assayed spectrophotometrically (at 540 nm for the example). A standard curve can be constructed using known concentrations of aniline, which will form a chromophore with similar properties to 2-aminophenol glucuronide.

Glutathione S-transferase activity of DME is measured using a model substrate, such as 2,4-dinitro-1-chlorobenzene, which reacts with glutathione to form a product, 2,4-dinitrophenyl-glutathione, that has an absorbance maximum at 340 nm. It is important to note that GSTs have differing substrate specificities, and the model substrate should be selected based on the substrate preferences of the GST of interest. Assays are performed at ambient temperature and contain an aliquot of the enzyme in a suitable reaction buffer (for example, 1 mM glutathione, 1 mM dinitrochlorobenzene, 90 mM potassium phosphate buffer pH 6.5). Reactions are carried out in an optical cuvette, and the absorbance at 340 nm is measured. The rate of increase in absorbance is proportional to the enzyme activity in the assay.

N-acyltransferase activity of DME is measured using radiolabeled amino acid substrates and measuring radiolabel incorporation into conjugated products. Enzyme is incubated in a reaction buffer containing an unlabeled acyl-CoA compound and radiolabeled amino acid, and the radiolabeled acyl-conjugates are separated from the unreacted amino acid by extraction into n-butanol or other appropriate organic solvent. For example, Johnson, M.R. et al. (1990; J. Biol. Chem. 266:10227-10233) measured bile acid-CoA:amino acid N-acyltransferase activity by incubating the enzyme with cholyl-CoA and <sup>3</sup>H-glycine or <sup>3</sup>H-aurine, separating the tritiated cholate conjugate by extraction into n-butanol, and measuring the radioactivity in the extracted product by scintillation. Alternatively, N-acyltransferase activity is measured using the spectrophotometric determination of reduced CoA (CoASH) described below.

N-acetyltransferase activity of DME is measured using the transfer of radiolabel from [ $^{14}\text{C}$ ]acetyl-CoA to a substrate molecule (for example, see Deguchi, T. (1975) J. Neurochem. 24:1083-5). Alternatively, a spectrophotometric assay based on DTNB (5,5'-dithio-bis(2-nitrobenzoic acid; Ellman's reagent) reaction with CoASH may be used. Free thiol-containing CoASH is formed during N-acetyltransferase catalyzed transfer of an acetyl group to a substrate. CoASH is detected using the absorbance of DTNB conjugate at 412 nm (De Angelis, J. et al. (1997) J. Biol. Chem. 273:3045-3050). Enzyme activity is proportional to the rate of radioactivity incorporation into substrate, or the rate of absorbance increase in the spectrophotometric assay.

Protein arginine methyltransferase activity of DME is measured at 37 °C for various periods of time. S-adenosyl-L-[methyl- $^3\text{H}$ ]methionine ([ $^3\text{H}$ ]AdoMet; specific activity = 75 Ci/mmol; NEN Life Science Products) is used as the methyl-donor substrate. Useful methyl-accepting substrates include glutathione S-transferase fibrillarlin glycine-arginine domain fusion protein (GST-GAR), heterogeneous nuclear ribonucleoprotein (hnRNP), or hypomethylated proteins present in lysates from adenosine dialdehyde-treated cells. Methylation reactions are stopped by adding SDS-PAGE sample buffer. The products of the reactions are resolved by SDS-PAGE and visualized by fluorography. The presence of  $^3\text{H}$ -labeled methyl-donor substrates is indicative of protein arginine methyltransferase activity of DME (Tang, J. et al. (2000) J. Biol. Chem. 275:7723-7730 and Tang, J. et al. (2000) J. Biol. Chem. 275:19866-19876).

Catechol-*O*-methyltransferase activity of DME is measured in a reaction mixture consisting of 50 mM Tris-HCl (pH 7.4), 1.2 mM  $\text{MgCl}_2$ , 200  $\mu\text{M}$  SAM (S-adenosyl-L-methionine) iodide (containing 0.5  $\mu\text{Ci}$  of methyl-[ $^3\text{H}$ ]SAM), 1 mM dithiothreitol, and varying concentrations of catechol substrate (e.g., L-dopa, dopamine, or DBA) in a final volume of 1.0 ml. The reaction is initiated by the addition of 250-500  $\mu\text{g}$  of purified DME or crude DME-containing sample and performed at 37 °C for 30 min. The reaction is arrested by rapidly cooling on ice and immediately extracting with 7 ml of ice-cold n-heptane. Following centrifugation at 1000 x g for 10 min, 3-ml aliquots of the organic extracts are analyzed for radioactivity content by liquid scintillation counting. The level of catechol-associated radioactivity in the organic phase is proportional to the catechol-*O*-methyltransferase activity of DME (Zhu, B.T. and J.G. Liehr (1996) 271:1357-1363).

DHFR activity of DME is determined spectrophotometrically at 15 °C by following the disappearance of NADPH at 340 nm ( $\epsilon_{340} = 11,800 \text{ M}^{-1}\text{cm}^{-1}$ ). The standard assay mixture contains 100  $\mu\text{M}$  NADPH, 14 mM 2-mercaptoethanol, MTEN buffer (50 mM 2-morpholinoethanesulfonic acid, 25 mM tris(hydroxymethyl)aminomethane, 25 mM ethanolamine, and 100 mM NaCl, pH 7.0), and DME in a final volume of 2.0 ml. The reaction is started by the addition of 50  $\mu\text{M}$  dihydrofolate (as substrate). The oxidation of NADPH to  $\text{NADP}^+$  corresponds to the reduction of dihydrofolate in the reaction and is proportional to the amount of DHFR activity in the sample (Nakamura, T. and



Iwakura, M. (1999) *J. Biol. Chem.* 274:19041-19047).

Aldo/keto reductase activity of DME is measured using the decrease in absorbance at 340 nm as NADPH is consumed. A standard reaction mixture is 135 mM sodium phosphate buffer (pH 6.2-7.2 depending on enzyme), 0.2 mM NADPH, 0.3 M lithium sulfate, 0.5-2.5 mg enzyme and an appropriate level of substrate. The reaction is incubated at 30°C and the reaction is monitored continuously with a spectrophotometer. Enzyme activity is calculated as mol NADPH consumed / mg of enzyme.

Alcohol dehydrogenase activity of DME is measured using the increase in absorbance at 340 nm as NAD<sup>+</sup> is reduced to NADH. A standard reaction mixture is 50 mM sodium phosphate, pH 7.5, and 0.25 mM EDTA. The reaction is incubated at 25°C and monitored using a spectrophotometer. Enzyme activity is calculated as mol NADH produced / mg of enzyme.

Carboxylesterase activity of DME activity is determined using 4-methylumbelliferyl acetate as a substrate. The enzymatic reaction is initiated by adding approximately 10 µl of DME-containing sample to 1 ml of reaction buffer (90 mM KH<sub>2</sub>PO<sub>4</sub>, 40 mM KCl, pH 7.3) with 0.5 mM 4-methylumbelliferyl acetate at 37°C. The production of 4-methylumbelliferone is monitored with a spectrophotometer ( $\epsilon_{350} = 12.2 \text{ mM}^{-1} \text{ cm}^{-1}$ ) for 1.5 min. Specific activity is expressed as micromoles of product formed per minute per milligram of protein and corresponds to the activity of DME in the sample (Evgenia, V. et al. (1997) *J. Biol. Chem.* 272:14769-14775).

In the alternative, the cocaine benzoyl ester hydrolase activity of DME is measured by incubating approximately 0.1 ml of enzyme and 3.3 mM cocaine in reaction buffer (50 mM NaH<sub>2</sub>PO<sub>4</sub>, pH 7.4) with 1 mM benzamidine, 1 mM EDTA, and 1 mM dithiothreitol at 37°C. The reaction is incubated for 1 h in a total volume of 0.4 ml then terminated with an equal volume of 5% trichloroacetic acid. 0.1 ml of the internal standard 3,4-dimethylbenzoic acid (10 µg/ml) is added. Precipitated protein is separated by centrifugation at 12,000 × g for 10 min. The supernatant is transferred to a clean tube and extracted twice with 0.4 ml of methylene chloride. The two extracts are combined and dried under a stream of nitrogen. The residue is resuspended in 14% acetonitrile, 250 mM KH<sub>2</sub>PO<sub>4</sub>, pH 4.0, with 8 µl of diethylamine per 100 ml and injected onto a C18 reverse-phase HPLC column for separation. The column eluate is monitored at 235 nm. DME activity is quantified by comparing peak area ratios of the analyte to the internal standard. A standard curve is generated with benzoic acid standards prepared in a trichloroacetic acid-treated protein matrix (Evgenia, V. et al. (1997) *J. Biol. Chem.* 272:14769-14775).

In another alternative, DME carboxyl esterase activity against the water-soluble substrate para-nitrophenyl butyric acid is determined by spectrophotometric methods well known to those skilled in the art. In this procedure, the DME-containing samples are diluted with 0.5 M Tris-HCl (pH 7.4 or 8.0) or sodium acetate (pH 5.0) in the presence of 6 mM taurocholate. The assay is

initiated by adding a freshly prepared para-nitrophenyl butyric acid solution (100 µg/ml in sodium acetate, pH 5.0). Carboxyl esterase activity is then monitored and compared with control autohydrolysis of the substrate using a spectrophotometer set at 405 nm (Wan, L. et al. (2000) J. Biol. Chem. 275:10041-10046).

5           Sulfotransferase activity of DME is measured using the incorporation of  $^{35}\text{S}$  from [ $^{35}\text{S}$ ]PAPS into a model substrate such as phenol (Folds, A. and J.L. Meek (1973) Biochim. Biophys. Acta 327:365-374). An aliquot of enzyme is incubated at 37°C with 1 mL of 10 mM phosphate buffer, pH 6.4, 50 mM phenol, and 0.4-4.0 mM [ $^{35}\text{S}$ ]PAPS. After sufficient time for 5-20% of the radiolabel to be transferred to the substrate, 0.2 mL of 0.1 M barium acetate is added to precipitate protein and  
10   phosphate buffer. Then 0.2 mL of 0.1 M  $\text{Ba}(\text{OH})_2$  is added, followed by 0.2 mL  $\text{ZnSO}_4$ . The supernatant is cleared by centrifugation, which removes proteins as well as unreacted [ $^{35}\text{S}$ ]PAPS. Radioactivity in the supernatant is measured by scintillation. The enzyme activity is determined from the number of moles of radioactivity in the reaction product.

          Heparan sulfate 6-sulfotransferase activity of DME is measured in vitro by incubating a  
15   sample containing DME along with 2.5 µmol imidazole HCl (pH 6.8), 3.75 µg of protamine chloride, 25 nmol (as hexosamine) of completely desulfated and N-resulfated heparin, and 50 pmol (about  $5 \times 10^5$  cpm) of [ $^{35}\text{S}$ ] adenosine 3'-phosphate 5'-phosphosulfate (PAPS) in a final reaction volume of 50 µl at 37°C for 20 min. The reaction is stopped by immersing the reaction tubes in a boiling water bath for 1 min. 0.1 µmol (as glucuronic acid) of chondroitin sulfate A is added to the reaction  
20   mixture as a carrier.  $^{35}\text{S}$ -Labeled polysaccharides are precipitated with 3 volumes of cold ethanol containing 1.3% potassium acetate and separated completely from unincorporated [ $^{35}\text{S}$ ]PAPS and its degradation products by gel chromatography using desalting columns. One unit of enzyme activity is defined as the amount required to transfer 1 pmol of sulfate/min., determined by the amount of [ $^{35}\text{S}$ ]PAPS incorporated into the precipitated polysaccharides (Habuchi, H. et al. (1995) J. Biol.  
25   Chem. 270:4172-4179).

          In the alternative, heparan sulfate 6-sulfotransferase activity of DME is measured by extraction and renaturation of enzyme from gels following separation by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). Following separation, the gel is washed with buffer (0.05 M Tris-HCl, pH 8.0), cut into 3-5 mm segments and subjected to agitation at 4 °C with  
30   100 µl of the same buffer containing 0.15 M NaCl for 48 h. The eluted enzyme is collected by centrifugation and assayed for the sulfotransferase activity as described above (Habuchi, H. et al. (1995) J. Biol. Chem. 270:4172-4179).

          In another alternative, DME sulfotransferase activity is determined by measuring the transfer of [ $^{35}\text{S}$ ]sulfate from [ $^{35}\text{S}$ ]PAPS to an immobilized peptide that represents the N-terminal 15 residues  
35   of the mature P-selectin glycoprotein ligand-1 polypeptide to which a C-terminal cysteine residue is

added. The peptide spans three potential tyrosine sulfation sites. The peptide is linked via the cysteine residue to iodoacetamide-activated resin at a density of 1.5-3.0  $\mu\text{mol}$  peptide/ml of resin. The enzyme assay is performed by combining 10  $\mu\text{l}$  of peptide-derivitized beads with 2-20  $\mu\text{l}$  of DME-containing sample in 40 mM Pipes (pH 6.8), 0.3 M NaCl, 20 mM  $\text{MnCl}_2$ , 50 mM NaF, 1% Triton X-100, and 1 mM 5'-AMP in a final volume of 130  $\mu\text{l}$ . The assay is initiated by addition of 0.5  $\mu\text{Ci}$  of [ $^{35}\text{S}$ ]PAPS (1.7  $\mu\text{M}$ ; 1 Ci = 37 GBq). After 30 min at 37°C, the reaction beads are washed with 6 M guanidine at 65°C and the radioactivity incorporated into the beads is determined by liquid scintillation counting. Transfer of [ $^{35}\text{S}$ ]sulfate to the bead-associated peptide is measured to determine the DME activity in the sample. One unit of activity is defined as 1 pmol of product formed per min (Ouyang, Y.-B. et al. (1998) *Biochemistry* 95:2896-2901).

In another alternative, DME sulfotransferase assays are performed using [ $^{35}\text{S}$ ]PAPS as the sulfate donor in a final volume of 30  $\mu\text{l}$ , containing 50 mM Hepes-NaOH (pH 7.0), 250 mM sucrose, 1 mM dithiothreitol, 14  $\mu\text{M}$  [ $^{35}\text{S}$ ]PAPS (15 Ci/mmol), and dopamine (25  $\mu\text{M}$ ), *p*-nitrophenol (5  $\mu\text{M}$ ), or other candidate substrates. Assay reactions are started by the addition of a purified DME enzyme preparation or a sample containing DME activity, allowed to proceed for 15 min at 37°C, and terminated by heating at 100°C for 3 min. The precipitates formed are cleared by centrifugation. The supernatants are then subjected to the analysis of  $^{35}\text{S}$ -sulfated product by either thin-layer chromatography or a two-dimensional thin layer separation procedure. Appropriate standards are run in parallel with the supernatants to allow the identification of the  $^{35}\text{S}$ -sulfated products and determine the enzyme specificity of the DME-containing samples based on relative rates of migration of reaction products (Sakakibara, Y. et al. (1998) *J. Biol. Chem.* 273:6242-6247).

Squalene epoxidase activity of DME is assayed in a mixture comprising purified DME (or a crude mixture comprising DME), 20 mM Tris-HCl (pH 7.5), 0.01 mM FAD, 0.2 unit of NADPH-cytochrome C (P-450) reductase, 0.01 mM [ $^{14}\text{C}$ ]squalene (dispersed with the aid of 20  $\mu\text{l}$  of Tween 80), and 0.2% Triton X-100. 1 mM NADPH is added to initiate the reaction followed by incubation at 37 °C for 30 min. The nonsaponifiable lipids are analyzed by silica gel TLC developed with ethyl acetate/benzene (0.5:99.5, v/v). The reaction products are compared to those from a reaction mixture without DME. The presence of 2,3(*S*)-oxidosqualene is confirmed using appropriate lipid standards (Sakakibara, J. et al. (1995) 270:17-20).

Epoxide hydrolase activity of DME is determined by following substrate depletion using gas chromatographic (GC) analysis of ethereal extracts or by following substrate depletion and diol production by GC analysis of reaction mixtures quenched in acetone. A sample containing DME or an epoxide hydrolase control sample is incubated in 10 mM Tris-HCl (pH 8.0), 1 mM ethylenediaminetetraacetate (EDTA), and 5 mM epoxide substrate (e.g., ethylene oxide, styrene oxide, propylene oxide, isoprene monoxide, epichlorohydrin, epibromohydrin, epifluorohydrin,

glycidol, 1,2-epoxybutane, 1,2-epoxyhexane, or 1,2-epoxyoctane). A portion of the sample is withdrawn from the reaction mixture at various time points, and added to 1 ml of ice-cold acetone containing an internal standard for GC analysis (e.g., 1-nonanol). Protein and salts are removed by centrifugation (15 min, 4000 × g) and the extract is analyzed by GC using a 0.2 mm × 25-m

5 CP-Wax57-CB column (CHROMPACK, Middelburg, The Netherlands) and a flame-ionization detector. The identification of GC products is performed using appropriate standards and controls well known to those skilled in the art. 1 Unit of DME activity is defined as the amount of enzyme that catalyzes the production of 1 μmol of diol/min (Rink, R. et al. (1997) *J. Biol. Chem.* 272:14650-14657).

10 Aminotransferase activity of DME is assayed by incubating samples containing DME for 1 hour at 37°C in the presence of 1 mM L-kynurenine and 1 mM 2-oxoglutarate in a final volume of 200 μl of 150 mM Tris acetate buffer (pH 8.0) containing 70 μM PLP. The formation of kynurenic acid is quantified by HPLC with spectrophotometric detection at 330 nm using the appropriate standards and controls well known to those skilled in the art. In the alternative,

15 L-3-hydroxykynurenine is used as substrate and the production of xanthurenic acid is determined by HPLC analysis of the products with UV detection at 340 nm. The production of kynurenic acid and xanthurenic acid, respectively, is indicative of aminotransferase activity (Buchli, R. et al. (1995) *J. Biol. Chem.* 270:29330-29335).

In another alternative, aminotransferase activity of DME is measured by determining the

20 activity of purified DME or crude samples containing DME toward various amino and oxo acid substrates under single turnover conditions by monitoring the changes in the UV/VIS absorption spectrum of the enzyme-bound cofactor, pyridoxal 5'-phosphate (PLP). The reactions are performed at 25°C in 50 mM 4-methylmorpholine (pH 7.5) containing 9 μM purified DME or DME containing samples and substrate to be tested (amino and oxo acid substrates). The half-reaction from amino

25 acid to oxo acid is followed by measuring the decrease in absorbance at 360 nm and the increase in absorbance at 330 nm due to the conversion of enzyme-bound PLP to pyridoxamine 5' phosphate (PMP). The specificity and relative activity of DME is determined by the activity of the enzyme preparation against specific substrates (Vacca, R.A. et al. (1997) *J. Biol. Chem.* 272:21932-21937).

Superoxide dismutase activity of DME is assayed from cell pellets, culture supernatants, or

30 purified protein preparations. Samples or lysates are resolved by electrophoresis on 15% non-denaturing polyacrylamide gels. The gels are incubated for 30 min in 2.5 mM nitro blue tetrazolium, followed by incubation for 20 min in 30 mM potassium phosphate, 30 mM TEMED, and 30 μM riboflavin (pH 7.8). Superoxide dismutase activity is visualized as white bands against a blue background, following illumination of the gels on a lightbox. Quantitation of superoxide

35 dismutase activity is performed by densitometric scanning of the activity gels using the appropriate

superoxide dismutase positive and negative controls (e.g., various amounts of commercially available E. coli superoxide dismutase (Harth, G. and Horwitz, M.A. (1999) J. Biol. Chem. 274:4281-4292).

#### **XVIII. Identification of DME Inhibitors**

5           Compounds to be tested are arrayed in the wells of a multi-well plate in varying concentrations along with an appropriate buffer and substrate, as described in the assays in Example XVII. DME activity is measured for each well and the ability of each compound to inhibit DME activity can be determined, as well as the dose-response profiles. This assay could also be used to identify molecules which enhance DME activity.

10           Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific  
15   embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID
2434655	1	2434655CD1	11	2434655CB1
2516747	2	2516747CD1	12	2516747CB1
7472775	3	7472775CD1	13	7472775CB1
7473323	4	7473323CD1	14	7473323CB1
7472777	5	7472777CD1	15	7472777CB1
1272843	6	1272843CD1	16	1272843CB1
7472790	7	7472790CD1	17	7472790CB1
7473944	8	7473944CD1	18	7473944CB1
2244136	9	2244136CD1	19	2244136CB1
7474327	10	7474327CD1	20	7474327CB1

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability Score	GenBank Homolog
1	2434655CD1	g5802604	4.70E-210	UDP glucuronosyltransferase UGT2A3 [Cavia porcellus] (Smith, S.A. et al. (1999) Mol. Genet. Metab. 68:68-77)
2	2516747CD1	g1246787	1.50E-69	UDP-galactose ceramide galactosyltransferase [Mus musculus] (Bosio, A. et al. (1996) Genomics 35:223-226)
3	7472775CD1	g2660716	2.10E-40	HNK-1 sulfotransferase [Rattus norvegicus] (Bakker, H. et al. (1997) J. Biol. Chem. 272:29942-29946)
		g12711481	0	N-acetylglactosamine 4-O-sulfotransferase 2 GalNAc4ST-2 [Homo sapiens]
4	7473323CD1	g2660716	1.50E-37	HNK-1 sulfotransferase [Rattus norvegicus] (Bakker, H. et al. (1997) J. Biol. Chem. 272:29942-29946)
		g11990885	0	GalNAc 4-sulfotransferase [Homo sapiens] (Okuda, T. et al. (2000) J. Biol. Chem. 275:40605-40613)
5	7472777CD1	g5917706	3.70E-178	N-acetylglucosamine 6-O-sulfotransferase [Homo sapiens] (Lee, J.K. et al. (1999) Biochem. Biophys. Res. Commun. 263:543-549)
6	1272843CD1	g402843	2.70E-134	Cytochrome P450 2B-Bx, phenobarbital-inducible [Oryctolagus cuniculus] (Ryan, R. et al. (1993) Arch. Biochem. Biophys. 304:454-463)
7	7472790CD1	g12653085	1.00E-104	N-acetyltransferase, homolog of S. cerevisiae ARD1 [Homo sapiens]
		g1302661	3.40E-97	ARD1 N-acetyl transferase related protein [Homo sapiens] (Brenner, V. et al. (1997) Genomics 44:8-14)
8	7473944CD1	g7340847	0	Chondroitin 4-sulfotransferase [Mus musculus] Yamauchi, S. et al. (2000) J. Biol. Chem. 275:8975-8981)
		g2921306	3.10E-39	HNK-1 sulfotransferase [Homo sapiens]
9	2244136CD1	g6651438	6.50E-23	Putative N-acetyltransferase Camello 2 [Homo sapiens]
10	7474327CD1	g178489	7.80E-153	Aldose reductase [Homo sapiens] (Chung, S. and S. LaMendola (1989) J. Biol. Chem. 264:14775-14777)

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	2434655CD1	527	S3 T120 T169 T200 S435 T520 S70 S343 S62 T141 T204 T243 T258 S296 S297 T419 Y130 Y143	N131 N313 N518	Signal peptide: M1-C23 Signal cleavage: M1-C23 Transmembrane domains: V91-N111, I492-F512 UDP-glucoronosyltransferase: G24-K525 UDP-glycosyltransferases BL00375: A: S34-L56; B: C125-P165; C: P188-N211; D: I253-C280; E: F293-P342; F: N348-P392; G: H447-H486 UDP-glucoronosyl and UDP-glucosyl transferases signature: N376-T417 UDP-glucoronosyltransferase (PD000190): G24-K338, V386-E527 UDP-glucoronosyl and glucosyl transferases: DM00367 P36510 176-459; M178-F460 UDP-glucoronosyltransferase signature: W354-Q397 Signal peptide: M1-A22 Signal cleavage: M1-L19 transmembrane domain: Y484-C503 UDP-glucoronosyl and UDP-glucosyl transferase: A23-K521 UDP-glucoronosyl and UDP-glucosyl transferases signature: N373-T416 UDP-glycosyltransferases BL00375: A: S34-L56; B: C121-P161; C: P180-N203; D: L246-E273; F: N345-P389; G: L444-Q483 UDP-glucoronosyltransferase: PD000190: K24-G291, L246-D430, L422-K522 UDP-glucoronosyl and UDP-glucosyl transferases: DM00367 P36513 188-462; L177-V456	HMMER SPScan HMMER HMMER-PFAM BLIMPS-BLOCKS ProfileScan BLAST-PRODOM BLAST-DOMO MOTIFS HMMER SPScan HMMER HMMER-PFAM ProfileScan BLIMPS-BLOCKS BLAST-PRODOM BLAST-DOMO
2	2516747CD1	523	S107 S81 S95 T99 S126 S213 S219 T224 S374 S59 S133 S362 S409 T423 Y74 Y433	N52		



Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
3	7472775CD1	358	S95 S42 S16 S19 T287 S315 T333 T44 S255 T335 Y251	N74 N158 N239 N352	HNK1 Sulfotransferase: PD041629; N65-L358	BLAST-PRODOM
4	7473323CD1	396	T44 S143 S144 T149 T356 S124 S190 S322 Y286 Y380	N100 N266 N339 N387	HNK1 Sulfotransferase: PD041629; L86-P391	BLAST-PRODOM
5	7472777CD1	395	S48 T214 T257 S322 T231 S363	N116 N229 N305 N328	Signal peptide: M1-A35 Signal cleavage: M1-S32 Sulfotransferase PD042460: V242-V375	HMMER SPScan BLAST-PRODOM
6	1272843CD1	504	T131 S119 S192 S343 S61 S99 T248 T288 T378 S473		Cytochrome P450: DM00022 S35666 49-475; L53-G480 Cytochrome P450: P33-R493 Cytochrome P450 cysteine heme-iron ligand signature: F412-S462 Cytochrome P450 PD000021: P33-Q150 Cytochrome P450: F433-G442 E-class P450 group I, PR00463: A: S61-V80; B: A86-L107; C: A177-D195; D: N291-S308; E: V311-G337; F: D354-P372; G: L395-D419; H: F430-C440; I: C440-L463 E-class P450 group IV, PR00465: B: S55-V78; C: L293-M319; D: L350-L366; E: F482-D496; F: L400-L418; G: F424-C440; H: C440-L458 P450 superfamily signature PR00385: A: A302-M319; B: K320-N333; C: A355-L366; D: L431-C440; E: C440-F451 Signal peptide: M1-A28 Signal cleavage: M1-A28 Transmembrane domains: W7-T26; I297-L315	BLAST-DOMO HMMER-PFAM ProfileScan BLAST-PRODOM MOTIFS BLIMPS-PRINTS BLIMPS-PRINTS BLIMPS-PRINTS HMMER SPScan HMMER

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
7	7472790CD1	229	S80 S114 S131 T184 S186 S202		Acetyltransferase (GNAT) family: I3-Q129 Acetyltransferase: PD071691: I130-E179 Acetyltransferase: DM04629 P41227 1-193: M1-C192	HMME-PFAM BLAST- PRODOM BLAST-DOMO
8	7473944CD1	347	T83 T90 S91 T134 T196 T244 S302 T98 S163 T244 T276 T305 S51 Y238	N200 N218 N316 N337	Signal cleavage: M1-R38 Transmembrane domain: C15-I30 Sulfotransferase: PD041629: L72-L344	SPScan HMME BLAST- PRODOM
9	2244136CD1	218	S69 S197	N156	Signal cleavage: M1-G63 Acetyltransferase: L43-R181 Aldo/keto reductase family: L7-R299	SPScan HMME-PFAM HMME-PFAM
10	7474327CD1	318	T76 S115 T240 S308 S216 S229	N9 N244	Aldo/keto reductase family: BL00798A: L7-W21 BL00798B: A35-I59 BL00798C: V68-W80 BL00798D: A92-Y108 BL00798E: K179-S216 BL00798F: T240-M288 Aldo-keto reductase signature: PR00069A: A35-I59, PR00069B: K95-P113, PR00069C: M147-F164, PR00069D: V183-Y212, PR00069E: L230-F254 Aldose reductase: PD000288: V28-N297 Aldo/keto reductase family: DM00192 P07943 1-297: A2-C301 Aldoketo reductase 2: M147-F164 Aldo/keto reductase family signatures: P124-Q186, I236-R299	BLIMPS- BLOCKS           BLAST- PRODOM BLAST-DOMO Motifs ProfileScan

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragments	Sequence Fragments	5' Position	3' Position
11	2434655CB1	1636	1-192, 196-294, 1531-1636, 1271-1328	71447910V1	400	1040
				3700065F6 (SININOT05)	1	615
				2096523R6 (BRAITUT02)	1257	1636
				1500810F6 (SINTBST01)	879	1429
				SXBC00120V1	1545	2086
12	2516747CB1	2086	964-1669, 2039-2086	SXBC01648V1	1189	1753
				SXBC00083V1	1040	1610
				SCSA02735V1	782	1141
				SXBC01451V1	1	544
				SXBC01528V1	283	863
13	7472775CB1	1814	1795-1814	2859356F6 (SININOT03)	611	1039
				3549975H1 (BRONDIT01)	170	416
				GNN:g6094626_000014_002:1-78,79- 1170	94	1185
				4189221H1 (BRSTNOT31)	517	845
				g2077168	1526	1814
14	7473323CB1	1650	1-664	60208402U1	357	812
				4166935T6 (PANCNOT21)	1151	1797
				g5663459	1395	1806
				g880305	1	513
				g6699998	827	1283
14	7473323CB1	1650	1-664	FL152824_00001	1	1637
				6197818H1 (PITUNON01)	1011	1650

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragments	Sequence Fragments	5' Position	3' Position
15	7472777CB1	1647	1-120, 802-1295	1593910F6 (BRAINT14)	1190	1647
				6869651H1 (BRAGNON02)	859	1301
				GNN.g7025744_000055_002	1	1188
16	1272843CB1	2620	1-552, 1122-1286, 1-1345	70792863V1	1150	1832
				5056523H1 (COLATMT01)	2413	2620
				1272843T6 (TESTTUT02)	1796	2594
				70567140V1	991	1560
				6576957H1 (COLHTUS02)	1836	2598
				6820526H1 (SINTNOR01)	209	831
17	7472790CB1	690	498-690	6823903H1 (SINTNOR01)	1	285
				6820526J1 (SINTNOR01)	377	1060
				GNN.g7024025_000010_002	1	690
				7391722H1 (LIVRFE02)	517	1113
				966072R1 (BRSTNOT05)	1015	1481
				6304381H2 (NERDTN03)	1181	1510
				7321512H1 (SPLNNOE01)	1	465
				6512789H1 (THYMDIT01)	315	922
				5961077F8 (BRATNOT05)	1259	1997
				7325258H1 (TESTTUE02)	395	997
19	2244136CB1	3701	3544-3701, 1-788, 2902-2953, 694-1425, 1677-1926, 2902-3701	7170810H2 (BRSTTMC01)	3023	3701
				5973836H1 (BRAZNOT01)	2649	3198
				7074871H1 (BRAUTDR04)	2145	2755
				1699688F6 (BLADTUT05)	1732	2352
				6637768H1 (BRACDIR02)	666	1352
				6888956J1 (BRAITDR03)	1	649
				GNN.g7407895_000004_002	1	960
20	7474327CB1	960	1-22			

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
11	2434655CB1	SINITMC01
12	2516747CB1	LIVRTUT04
13	7472775CB1	PANCNOT21
14	7473323CB1	PITUNON01
15	7472777CB1	BRAINOT14
16	1272843CB1	SINTNOR01
18	7473944CB1	BRAINOT09
19	2244136CB1	BRAITDR03

Table 6

Library	Vector	Library Description
BRAINOT09	pINCY	Library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus, who died at 23 weeks' gestation.
BRAINOT14	pINCY	Library was constructed using RNA isolated from brain tissue removed from the left frontal lobe of a 40-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology for the associated tumor tissue indicated grade 4 gemistocytic astrocytoma.
BRAITDR03	PCDNA2.1	This random primed library was constructed using RNA isolated from allocortex, cingulate posterior tissue removed from a 55-year-old Caucasian female who died from cholangiocarcinoma. Pathology indicated mild meningeal fibrosis predominately over the convexities, scattered axonal spheroids in the white matter of the cingulate cortex and the thalamus, and a few scattered neurofibrillary tangles in the entorhinal cortex and the periaqueductal gray region. Pathology for the associated tumor tissue indicated well-differentiated cholangiocarcinoma of the liver with residual or relapsed tumor. Patient history included cholangiocarcinoma, post-operative Budd-Chiari syndrome, biliary ascites, hydorthorax, dehydration, malnutrition, oliguria and acute renal failure. Previous surgeries included cholecystectomy and resection of 85% of the liver.
LIVRTUT04	pINCY	Library was constructed using RNA isolated from liver tumor tissue removed from a 50-year-old Caucasian male during a partial hepatectomy. Pathology indicated a grade 3-4 hepatoma, forming a mass. Patient history included benign hypertension and hepatitis. Hepatitis B core antigen and hepatitis B surface antigen was present in the patient.
PANCN0T21	pINCY	Library was constructed using RNA isolated from pancreatic tissue removed from an 8-year-old Black male who died from anoxia.
SINTNOR01	PCDNA2.1	This random primed library was constructed using RNA isolated from small intestine tissue removed from a 31-year-old Caucasian female during Roux-en-Y gastric bypass. Patient history included clinical obesity.

Table 6 (cont.)

Library	Vector	Library Description
PITUNON01	pINCY	<p>This normalized pituitary gland tissue library was constructed from 6.92 million independent clones from a pituitary gland tissue library. Starting RNA was made from pituitary gland tissue removed from a 55-year-old male who died from chronic obstructive pulmonary disease. Neuropathology indicated there were no gross abnormalities, other than mild ventricular enlargement. There was no apparent microscopic abnormality in any of the neocortical areas examined, except for a number of silver positive neurons with apical dendrite staining, particularly in the frontal lobe. The significance of this was undetermined. The only other microscopic abnormality was that there was prominent silver staining with some swollen axons in the CA3 region of the anterior and posterior hippocampus. Microscopic sections of the cerebellum revealed mild Bergmann's gliosis in the Purkinje cell layer. Patient history included schizophrania. The library was normalized in two rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228-9232 and Bonaldo et al., Genome Research (1996) 6:791, except that a significantly longer (48 hours/round) reannealing hybridization was used.</p>
SINITMC01	pINCY	<p>This large size-fractionated library was constructed using pooled cDNA from two donors. cDNA was generated using mRNA isolated from ileum tissue removed from a 30-year-old Caucasian female (donor A) during partial colectomy, open liver biopsy, and permanent colostomy, and from ileum tissue removed from a 70-year-old Caucasian female (donor B) during right hemicolectomy, open liver biopsy, sigmoidoscopy, colonoscopy, and permanent colostomy. Pathology for the matched tumor tissue (donor A) indicated carcinoid tumor (grade 1 neuroendocrine carcinoma) arising in the terminal ileum. The tumor permeated through the ileal wall into the mesenteric fat and extended into the adherent cecum, where tumor extended through the bowel wall up to the mucosal surface. Multiple lymph nodes were positive for tumor. Additional (2) lymph nodes were also involved by direct tumor extension. Pathology for donor B indicated a non-tumorous margin of ileum. Pathology for the matched tumor (donor B) indicated invasive grade 2 adenocarcinoma forming an ulcerated mass, situated distal to the ileocecal valve. The tumor invaded through the muscularis propria just into the serosal adipose tissue. One regional lymph node was positive for a microfocus of metastatic adenocarcinoma. Donor A presented with flushing and unspecified abdominal/pelvic symptoms. Patient history included endometriosis, and tobacco and alcohol abuse. Donor B's history included a malignant breast neoplasm, type II diabetes, hyperlipidemia, viral hepatitis, an unspecified thyroid disorder, osteoarthritis, and a malignant skin neoplasm. Donor B's medication included tamoxifen.</p>

Table 7

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322; Durbin, R. et al. (1998) Our World View, in a Nutshell, Cambridge Univ. Press, pp. 1-350.	PFAM hits: Probability value= 1.0E-3 or less Signal peptide hits: Score= 0 or greater



Table 7 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score $\geq$ GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:

a) a polypeptide comprising an amino acid sequence selected from the group consisting of

5 SEQ ID NO:1-10,

b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-10,

c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, and

10 d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10.

2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-10.

15

3. An isolated polynucleotide encoding a polypeptide of claim 1.

4. An isolated polynucleotide encoding a polypeptide of claim 2.

20 5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:11-20.

6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

25

7. A cell transformed with a recombinant polynucleotide of claim 6.

8. A transgenic organism comprising a recombinant polynucleotide of claim 6.

30 9. A method for producing a polypeptide of claim 1, the method comprising:

a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

5 11. An isolated polynucleotide selected from the group consisting of:

a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:11-20,

b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:11-20,

10 c) a polynucleotide complementary to a polynucleotide of a),

d) a polynucleotide complementary to a polynucleotide of b), and

e) an RNA equivalent of a)-d).

12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a  
15 polynucleotide of claim 11.

13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

20 a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and

b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

25

14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

30 a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and

b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. A composition of claim 16, wherein the polypeptide has an amino acid sequence  
5 selected from the group consisting of SEQ ID NO:1-10.

18. A method for treating a disease or condition associated with decreased expression of functional DME, comprising administering to a patient in need of such treatment the composition of claim 16.

10 19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

15 20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

20 21. A method for treating a disease or condition associated with decreased expression of functional DME, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- 25
- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
  - b) detecting antagonist activity in the sample.

23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

30 24. A method for treating a disease or condition associated with overexpression of functional DME, comprising administering to a patient in need of such treatment a composition of claim 23.

25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and

5 b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

10 a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,

b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and

15 c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,

b) detecting altered expression of the target polynucleotide, and

25 c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

28. A method for assessing toxicity of a test compound, said method comprising:

a) treating a biological sample containing nucleic acids with the test compound;

30 b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;

35 c) quantifying the amount of hybridization complex; and

d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

5

29. A diagnostic test for a condition or disease associated with the expression of DME in a biological sample comprising the steps of:

a) combining the biological sample with an antibody of claim 10, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex; and

10

b) detecting the complex, wherein the presence of the complex correlates with the presence of the polypeptide in the biological sample.

30. The antibody of claim 10, wherein the antibody is:

a) a chimeric antibody,

15

b) a single chain antibody,

c) a Fab fragment,

d) a F(ab')<sub>2</sub> fragment, or

e) a humanized antibody.

20

31. A composition comprising an antibody of claim 10 and an acceptable excipient.

32. A method of diagnosing a condition or disease associated with the expression of DME in a subject, comprising administering to said subject an effective amount of the composition of claim 31.

25

33. A composition of claim 31, wherein the antibody is labeled.

34. A method of diagnosing a condition or disease associated with the expression of DME in a subject, comprising administering to said subject an effective amount of the composition of claim 33.

30

35. A method of preparing a polyclonal antibody with the specificity of the antibody of claim 10 comprising:

a) immunizing an animal with a polypeptide having an amino acid sequence selected from

the group consisting of SEQ ID NO:1-10, or an immunogenic fragment thereof, under conditions to elicit an antibody response;

b) isolating antibodies from said animal; and

5 c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10.

36. An antibody produced by a method of claim 35.

10 37. A composition comprising the antibody of claim 36 and a suitable carrier.

38. A method of making a monoclonal antibody with the specificity of the antibody of claim 10 comprising:

15 a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10, or an immunogenic fragment thereof, under conditions to elicit an antibody response;

b) isolating antibody producing cells from the animal;

c) fusing the antibody producing cells with immortalized cells to form monoclonal antibody-producing hybridoma cells;

20 d) culturing the hybridoma cells; and

e) isolating from the culture monoclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10.

25 39. A monoclonal antibody produced by a method of claim 38.

40. A composition comprising the antibody of claim 39 and a suitable carrier.

41. The antibody of claim 10, wherein the antibody is produced by screening a Fab expression library.

30 42. The antibody of claim 10, wherein the antibody is produced by screening a recombinant immunoglobulin library.

43. A method for detecting a polypeptide having an amino acid sequence selected from the

group consisting of SEQ ID NO:1-10 in a sample, comprising the steps of:

a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and

b) detecting specific binding, wherein specific binding indicates the presence of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10 in the sample.

44. A method of purifying a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10 from a sample, the method comprising:

a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and

b) separating the antibody from the sample and obtaining the purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-10.

45. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.

46. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2.

47. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:3.

48. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:4.

49. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:5.

50. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:6.

51. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:7.

52. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:8.

53. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:9.

54. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:10.



55. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:11.

56. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
5 NO:12.

57. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:13.

10 58. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:14.

59. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:15.  
15

60. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:16.

61. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
20 NO:17.

62. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:18.

25 63. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:19.

64. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID  
NO:20.

<110> INCYTE GENOMICS, INC.  
 POLICKY, Jennifer L.  
 HAFALIA, April  
 BURFORD, Neil  
 RING, Huijun Z.  
 LAL, Preeti  
 TRIBOULEY, Catherine M.  
 YAO, Monique G.  
 YUE, Henry  
 TANG, Y. Tom  
 PATTERSON, Chandra  
 DAS, Debopriya  
 SANJANWALA, Madhu S.  
 GANDHI, Ameena R.  
 REDDY, Roopa  
 KHAN, Farrah A.  
 BAUGHN, Mariah R.  
 RAMKUMAR, Jaya  
 GRIFFIN, Jennifer A.  
 AU-YOUNG, Janice

<120> DRUG METABOLIZING ENZYMES

<130> PI-0070 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/197,590; 60/198,403; 60/200,185; 60/202,234; 60/203,509

<151> 2000-04-13; 2000-04-19; 2000-04-28; 2000-05-05; 2000-05-11

<160> 20

<170> PERL Program

<210> 1

<211> 527

<212> PRT

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 2434655CD1

<400> 1

Met	Arg	Ser	Asp	Lys	Ser	Ala	Leu	Val	Phe	Leu	Leu	Leu	Gln	Leu
1				5					10					15
Phe	Cys	Val	Gly	Cys	Gly	Phe	Cys	Gly	Lys	Val	Leu	Val	Trp	Pro
			20						25					30
Cys	Asp	Met	Ser	His	Trp	Leu	Asn	Val	Lys	Val	Ile	Leu	Glu	Glu
				35					40					45
Leu	Ile	Val	Arg	Gly	His	Glu	Val	Thr	Val	Leu	Thr	His	Ser	Lys
				50					55					60
Pro	Ser	Leu	Ile	Asp	Tyr	Arg	Lys	Pro	Ser	Ala	Leu	Lys	Phe	Glu
				65					70					75
Val	Val	His	Met	Pro	Gln	Asp	Arg	Thr	Glu	Glu	Asn	Glu	Ile	Phe
				80					85					90
Val	Asp	Leu	Ala	Leu	Asn	Val	Leu	Pro	Gly	Leu	Ser	Thr	Trp	Gln
				95					100					105
Ser	Val	Ile	Lys	Leu	Asn	Asp	Phe	Phe	Val	Glu	Ile	Arg	Gly	Thr
				110					115					120
Leu	Lys	Met	Met	Cys	Glu	Ser	Phe	Ile	Tyr	Asn	Gln	Thr	Leu	Met
				125					130					135

Lys	Lys	Leu	Gln	Glu	Thr	Asn	Tyr	Asp	Val	Met	Leu	Ile	Asp	Pro
				140					145					150
Val	Ile	Pro	Cys	Gly	Asp	Leu	Met	Ala	Glu	Leu	Leu	Ala	Val	Pro
				155					160					165
Phe	Val	Leu	Thr	Leu	Arg	Ile	Ser	Val	Gly	Gly	Asn	Met	Glu	Arg
				170					175					180
Ser	Cys	Gly	Lys	Leu	Pro	Ala	Pro	Leu	Ser	Tyr	Val	Pro	Val	Pro
				185					190					195
Met	Thr	Gly	Leu	Thr	Asp	Arg	Met	Thr	Phe	Leu	Glu	Arg	Val	Lys
				200					205					210
Asn	Ser	Met	Leu	Ser	Val	Leu	Phe	His	Phe	Trp	Ile	Gln	Asp	Tyr
				215					220					225
Asp	Tyr	His	Phe	Trp	Glu	Glu	Phe	Tyr	Ser	Lys	Ala	Leu	Gly	Arg
				230					235					240
Pro	Thr	Thr	Leu	Cys	Glu	Thr	Val	Gly	Lys	Ala	Glu	Ile	Trp	Leu
				245					250					255
Ile	Arg	Thr	Tyr	Trp	Asp	Phe	Glu	Phe	Pro	Gln	Pro	Tyr	Gln	Pro
				260					265					270
Asn	Phe	Glu	Phe	Val	Gly	Gly	Leu	His	Cys	Lys	Pro	Ala	Lys	Ala
				275					280					285
Leu	Pro	Lys	Glu	Met	Glu	Asn	Phe	Val	Gln	Ser	Ser	Gly	Glu	Asp
				290					295					300
Gly	Ile	Val	Val	Phe	Ser	Leu	Gly	Ser	Leu	Phe	Gln	Asn	Val	Thr
				305					310					315
Glu	Glu	Lys	Ala	Asn	Ile	Ile	Ala	Ser	Ala	Leu	Ala	Gln	Ile	Pro
				320					325					330
Gln	Lys	Val	Leu	Trp	Arg	Tyr	Lys	Gly	Lys	Lys	Pro	Ser	Thr	Leu
				335					340					345
Gly	Ala	Asn	Thr	Arg	Leu	Tyr	Asp	Trp	Ile	Pro	Gln	Asn	Asp	Leu
				350					355					360
Leu	Gly	His	Pro	Lys	Thr	Lys	Ala	Phe	Ile	Thr	His	Gly	Gly	Met
				365					370					375
Asn	Gly	Ile	Tyr	Glu	Ala	Ile	Tyr	His	Gly	Val	Pro	Met	Val	Gly
				380					385					390
Val	Pro	Ile	Phe	Gly	Asp	Gln	Leu	Asp	Asn	Ile	Ala	His	Met	Lys
				395					400					405
Ala	Lys	Gly	Ala	Ala	Val	Glu	Ile	Asn	Phe	Lys	Thr	Met	Thr	Ser
				410					415					420
Glu	Asp	Leu	Leu	Arg	Ala	Leu	Arg	Thr	Val	Ile	Thr	Asp	Ser	Ser
				425					430					435
Tyr	Lys	Glu	Asn	Ala	Met	Arg	Leu	Ser	Arg	Ile	His	His	Asp	Gln
				440					445					450
Pro	Val	Lys	Pro	Leu	Asp	Arg	Ala	Val	Phe	Trp	Ile	Glu	Phe	Val
				455					460					465
Met	Arg	His	Lys	Gly	Ala	Lys	His	Leu	Arg	Ser	Ala	Ala	His	Asp
				470					475					480
Leu	Thr	Trp	Phe	Gln	His	Tyr	Ser	Ile	Asp	Val	Ile	Gly	Phe	Leu
				485					490					495
Leu	Thr	Cys	Val	Ala	Thr	Ala	Ile	Phe	Leu	Phe	Thr	Lys	Cys	Phe
				500					505					510
Leu	Phe	Ser	Cys	Gln	Lys	Phe	Asn	Lys	Thr	Arg	Lys	Ile	Glu	Lys
				515					520					525

Arg Glu

&lt;210&gt; 2

&lt;211&gt; 523

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2516747CD1

&lt;400&gt; 2

Met	Val	Gly	Gln	Arg	Val	Leu	Leu	Leu	Val	Ala	Phe	Leu	Leu	Ser
1				5					10					15
Gly	Val	Leu	Leu	Ser	Glu	Ala	Ala	Lys	Ile	Leu	Thr	Ile	Ser	Thr
				20					25					30
Leu	Gly	Gly	Ser	His	Tyr	Leu	Leu	Leu	Asp	Arg	Val	Ser	Gln	Ile
				35					40					45
Leu	Gln	Glu	His	Gly	His	Asn	Val	Thr	Met	Leu	His	Gln	Ser	Gly
				50					55					60
Lys	Phe	Leu	Ile	Pro	Asp	Ile	Lys	Glu	Glu	Glu	Lys	Ser	Tyr	Gln
				65					70					75
Val	Ile	Arg	Trp	Phe	Ser	Pro	Glu	Asp	His	Gln	Lys	Arg	Ile	Lys
				80					85					90
Lys	His	Phe	Asp	Ser	Tyr	Ile	Glu	Thr	Ala	Leu	Asp	Gly	Arg	Lys
				95					100					105
Glu	Ser	Glu	Ala	Leu	Val	Lys	Leu	Met	Glu	Ile	Phe	Gly	Thr	Gln
				110					115					120
Cys	Ser	Tyr	Leu	Leu	Ser	Arg	Lys	Asp	Ile	Met	Asp	Ser	Leu	Lys
				125					130					135
Asn	Glu	Asn	Tyr	Asp	Leu	Val	Phe	Val	Glu	Ala	Phe	Asp	Phe	Cys
				140					145					150
Ser	Phe	Leu	Ile	Ala	Glu	Lys	Leu	Val	Lys	Pro	Phe	Val	Ala	Ile
				155					160					165
Leu	Pro	Thr	Thr	Phe	Gly	Ser	Leu	Asp	Phe	Gly	Leu	Pro	Ser	Pro
				170					175					180
Leu	Ser	Tyr	Val	Pro	Val	Phe	Pro	Ser	Leu	Leu	Thr	Asp	His	Met
				185					190					195
Asp	Phe	Trp	Gly	Arg	Val	Lys	Asn	Phe	Leu	Met	Phe	Phe	Ser	Phe
				200					205					210
Ser	Arg	Ser	Gln	Trp	Asp	Met	Gln	Ser	Thr	Phe	Asp	Asn	Thr	Ile
				215					220					225
Lys	Glu	His	Phe	Pro	Glu	Gly	Ser	Arg	Pro	Val	Leu	Ser	His	Leu
				230					235					240
Leu	Leu	Lys	Ala	Glu	Leu	Trp	Phe	Val	Asn	Ser	Asp	Phe	Ala	Phe
				245					250					255
Asp	Phe	Ala	Arg	Pro	Leu	Leu	Pro	Asn	Thr	Val	Tyr	Ile	Gly	Gly
				260					265					270
Leu	Met	Glu	Lys	Pro	Ile	Lys	Pro	Val	Pro	Gln	Asp	Leu	Asp	Asn
				275					280					285
Phe	Ile	Ala	Asn	Phe	Gly	Asp	Ala	Gly	Phe	Val	Leu	Val	Ala	Phe
				290					295					300
Gly	Ser	Met	Leu	Asn	Thr	His	Gln	Ser	Gln	Glu	Val	Leu	Lys	Lys
				305					310					315
Met	His	Asn	Ala	Phe	Ala	His	Leu	Pro	Gln	Gly	Val	Ile	Trp	Thr
				320					325					330
Cys	Gln	Ser	Ser	His	Trp	Pro	Arg	Asp	Val	His	Leu	Ala	Thr	Asn
				335					340					345
Val	Lys	Ile	Val	Asp	Trp	Leu	Pro	Arg	Ser	Asp	Leu	Leu	Ala	His
				350					355					360
Pro	Ser	Ile	Arg	Leu	Phe	Val	Thr	His	Gly	Gly	Gln	Asn	Ser	Val
				365					370					375
Met	Glu	Ala	Ile	Arg	His	Gly	Val	Pro	Met	Val	Gly	Leu	Pro	Val
				380					385					390
Asn	Gly	Asp	Gln	His	Gly	Asn	Met	Val	Arg	Val	Val	Ala	Lys	Asn
				395					400					405
Tyr	Gly	Val	Ser	Ile	Arg	Leu	Asn	Gln	Val	Thr	Ala	Asp	Thr	Leu
				410					415					420
Thr	Leu	Thr	Met	Lys	Gln	Val	Ile	Glu	Asp	Lys	Arg	Tyr	Lys	Ser
				425					430					435
Ala	Val	Val	Ala	Ala	Ser	Val	Ile	Leu	His	Ser	Gln	Pro	Leu	Ser
				440					445					450
Pro	Ala	Gln	Arg	Leu	Val	Gly	Trp	Ile	Asp	His	Ile	Leu	Gln	Thr
				455					460					465

Gly	Gly	Ala	Thr	His	Leu	Lys	Pro	Tyr	Ala	Phe	Gln	Gln	Pro	Trp	
				470					475					480	
His	Glu	Gln	Tyr	Leu	Ile	Asp	Val	Phe	Val	Phe	Leu	Leu	Gly	Leu	
				485					490					495	
Thr	Leu	Gly	Thr	Met	Trp	Leu	Cys	Gly	Lys	Leu	Leu	Gly	Val	Val	
				500					505					510	
Ala	Arg	Trp	Leu	Arg	Gly	Ala	Arg	Lys	Val	Lys	Lys	Thr			
				515					520						

&lt;210&gt; 3

&lt;211&gt; 358

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7472775CD1

&lt;400&gt; 3

Met	Pro	Glu	Asp	Val	Arg	Glu	Lys	Lys	Glu	Asn	Leu	Leu	Leu	Asn	
1				5					10					15	
Ser	Glu	Arg	Ser	Thr	Arg	Leu	Leu	Thr	Lys	Thr	Ser	His	Ser	Gln	
				20					25					30	
Gly	Gly	Asp	Gln	Ala	Leu	Ser	Lys	Ser	Thr	Gly	Ser	Pro	Thr	Glu	
				35					40					45	
Lys	Leu	Ile	Glu	Lys	Arg	Gln	Gly	Ala	Lys	Thr	Val	Phe	Asn	Lys	
				50					55					60	
Phe	Ser	Asn	Met	Asn	Trp	Pro	Val	Asp	Ile	His	Pro	Leu	Asn	Lys	
				65					70					75	
Ser	Leu	Val	Lys	Asp	Asn	Lys	Trp	Lys	Lys	Thr	Glu	Glu	Thr	Gln	
				80					85					90	
Glu	Lys	Arg	Arg	Ser	Phe	Leu	Gln	Glu	Phe	Cys	Lys	Lys	Tyr	Gly	
				95					100					105	
Gly	Val	Ser	His	His	Gln	Ser	His	Leu	Phe	His	Thr	Val	Ser	Arg	
				110					115					120	
Ile	Tyr	Val	Glu	Asp	Lys	His	Lys	Ile	Leu	Tyr	Cys	Glu	Val	Pro	
				125					130					135	
Lys	Ala	Gly	Cys	Ser	Asn	Trp	Lys	Arg	Ile	Leu	Met	Val	Leu	Asn	
				140					145					150	
Gly	Leu	Ala	Ser	Ser	Ala	Tyr	Asn	Ile	Ser	His	Asn	Ala	Val	His	
				155					160					165	
Tyr	Gly	Lys	His	Leu	Lys	Lys	Leu	Asp	Ser	Phe	Asp	Leu	Lys	Gly	
				170					175					180	
Ile	Tyr	Thr	Arg	Leu	Asn	Thr	Tyr	Thr	Lys	Ala	Val	Phe	Val	Arg	
				185					190					195	
Asp	Pro	Met	Glu	Arg	Leu	Val	Ser	Ala	Phe	Arg	Asp	Lys	Phe	Glu	
				200					205					210	
His	Pro	Asn	Ser	Tyr	Tyr	His	Pro	Val	Phe	Gly	Lys	Ala	Ile	Ile	
				215					220					225	
Lys	Lys	Tyr	Arg	Pro	Asn	Ala	Cys	Glu	Glu	Ala	Leu	Ile	Asn	Gly	
				230					235					240	
Ser	Gly	Val	Lys	Phe	Lys	Glu	Phe	Ile	His	Tyr	Leu	Leu	Asp	Ser	
				245					250					255	
His	Arg	Pro	Val	Gly	Met	Asp	Ile	His	Trp	Glu	Lys	Val	Ser	Lys	
				260					265					270	
Leu	Cys	Tyr	Pro	Cys	Leu	Ile	Asn	Tyr	Asp	Phe	Val	Gly	Lys	Phe	
				275					280					285	
Glu	Thr	Leu	Glu	Glu	Asp	Ala	Asn	Tyr	Phe	Leu	Gln	Met	Ile	Gly	
				290					295					300	
Ala	Pro	Lys	Glu	Leu	Lys	Phe	Pro	Asn	Phe	Lys	Asp	Arg	His	Ser	
				305					310					315	
Ser	Asp	Glu	Arg	Thr	Asn	Ala	Gln	Val	Val	Arg	Gln	Tyr	Leu	Lys	
				320					325					330	

Asp	Leu	Thr	Arg	Thr	Glu	Arg	Gln	Leu	Ile	Tyr	Asp	Phe	Tyr	Tyr
				335					340					345
Leu	Asp	Tyr	Leu	Met	Phe	Asn	Tyr	Thr	Thr	Pro	Phe	Leu		
				350					355					

<210> 4  
 <211> 396  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 7473323CD1

<400> 4

Met	Ala	Ile	Asp	Ala	Leu	Val	Ser	Leu	Cys	Leu	Pro	Glu	Val	Ile
1				5					10					15
Arg	Ile	Lys	Phe	Asn	Ile	Arg	Pro	Arg	Gln	Pro	His	His	Asp	Leu
				20					25					30
Pro	Pro	Gly	Gly	Ser	Gln	Asp	Gly	Asp	Leu	Lys	Glu	Pro	Thr	Glu
				35					40					45
Arg	Val	Thr	Arg	Asp	Leu	Ser	Ser	Gly	Ala	Pro	Arg	Gly	Arg	Asn
				50					55					60
Leu	Pro	Ala	Pro	Asp	Gln	Pro	Gln	Pro	Pro	Leu	Gln	Arg	Gly	Thr
				65					70					75
Arg	Leu	Arg	Leu	Arg	Gln	Arg	Arg	Arg	Arg	Leu	Leu	Ile	Lys	Lys
				80					85					90
Met	Pro	Ala	Ala	Ala	Thr	Ile	Pro	Ala	Asn	Ser	Ser	Asp	Ala	Pro
				95					100					105
Phe	Ile	Arg	Pro	Gly	Pro	Gly	Thr	Leu	Asp	Gly	Arg	Trp	Val	Ser
				110					115					120
Leu	His	Arg	Ser	Gln	Gln	Glu	Arg	Lys	Arg	Val	Met	Gln	Glu	Ala
				125					130					135
Cys	Ala	Lys	Tyr	Arg	Ala	Ser	Ser	Ser	Arg	Arg	Ala	Val	Thr	Pro
				140					145					150
Arg	His	Val	Ser	Arg	Ile	Phe	Val	Glu	Asp	Arg	His	Arg	Val	Leu
				155					160					165
Tyr	Cys	Glu	Val	Pro	Lys	Ala	Gly	Cys	Ser	Asn	Trp	Lys	Arg	Val
				170					175					180
Leu	Met	Val	Leu	Ala	Gly	Leu	Ala	Ser	Ser	Thr	Ala	Asp	Ile	Gln
				185					190					195
His	Asn	Thr	Val	His	Tyr	Gly	Ser	Ala	Leu	Lys	Arg	Leu	Asp	Thr
				200					205					210
Phe	Asp	Arg	Gln	Gly	Ile	Leu	His	Arg	Leu	Ser	Thr	Tyr	Thr	Lys
				215					220					225
Met	Leu	Phe	Val	Arg	Glu	Pro	Phe	Glu	Arg	Leu	Val	Ser	Ala	Phe
				230					235					240
Arg	Asp	Lys	Phe	Glu	His	Pro	Asn	Ser	Tyr	Tyr	His	Pro	Val	Phe
				245					250					255
Gly	Lys	Ala	Ile	Leu	Ala	Arg	Tyr	Arg	Ala	Asn	Ala	Ser	Arg	Glu
				260					265					270
Ala	Leu	Arg	Thr	Gly	Ser	Gly	Val	Arg	Phe	Pro	Glu	Phe	Val	Gln
				275					280					285
Tyr	Leu	Leu	Asp	Val	His	Arg	Pro	Val	Gly	Met	Asp	Ile	His	Trp
				290					295					300
Asp	His	Val	Ser	Arg	Leu	Cys	Ser	Pro	Cys	Leu	Ile	Asp	Tyr	Asp
				305					310					315
Phe	Val	Gly	Lys	Phe	Glu	Ser	Met	Glu	Asp	Asp	Ala	Asn	Phe	Phe
				320					325					330
Leu	Ser	Leu	Ile	Arg	Ala	Pro	Arg	Asn	Leu	Thr	Phe	Pro	Arg	Phe
				335					340					345
Lys	Asp	Arg	His	Ser	Gln	Glu	Ala	Arg	Thr	Thr	Ala	Arg	Ile	Ala
				350					355					360

His	Gln	Tyr	Phe	Ala	Gln	Leu	Ser	Ala	Leu	Gln	Arg	Gln	Arg	Thr
				365					370					375
Tyr	Asp	Phe	Tyr	Tyr	Met	Asp	Tyr	Leu	Met	Phe	Asn	Tyr	Ser	Lys
				380					385					390
Pro	Phe	Ala	Asp	Leu	Tyr									
				395										

&lt;210&gt; 5

&lt;211&gt; 395

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7472777CD1

&lt;400&gt; 5

Met	Trp	Leu	Pro	Arg	Val	Ser	Ser	Thr	Ala	Val	Thr	Ala	Leu	Leu
1				5					10					15
Leu	Ala	Gln	Thr	Phe	Leu	Leu	Leu	Phe	Leu	Val	Ser	Arg	Pro	Gly
				20					25					30
Pro	Ser	Ser	Pro	Ala	Gly	Gly	Glu	Ala	Arg	Val	His	Val	Leu	Val
				35					40					45
Leu	Ser	Ser	Trp	Arg	Ser	Gly	Ser	Ser	Phe	Val	Gly	Gln	Leu	Phe
				50					55					60
Asn	Gln	His	Pro	Asp	Val	Phe	Tyr	Leu	Met	Glu	Pro	Ala	Trp	His
				65					70					75
Val	Trp	Thr	Thr	Leu	Ser	Gln	Gly	Ser	Ala	Ala	Thr	Leu	His	Met
				80					85					90
Ala	Val	Arg	Asp	Leu	Val	Arg	Ser	Val	Phe	Leu	Cys	Asp	Met	Asp
				95					100					105
Val	Phe	Asp	Ala	Tyr	Leu	Pro	Trp	Arg	Arg	Asn	Leu	Ser	Asp	Leu
				110					115					120
Phe	Gln	Trp	Ala	Val	Ser	Arg	Ala	Leu	Cys	Ser	Pro	Pro	Ala	Cys
				125					130					135
Ser	Ala	Phe	Pro	Arg	Gly	Ala	Ile	Ser	Ser	Glu	Ala	Val	Cys	Lys
				140					145					150
Pro	Leu	Cys	Ala	Arg	Gln	Ser	Phe	Thr	Leu	Ala	Arg	Glu	Ala	Cys
				155					160					165
Arg	Ser	Tyr	Ser	His	Val	Val	Leu	Lys	Glu	Val	Arg	Phe	Phe	Asn
				170					175					180
Leu	Gln	Val	Leu	Tyr	Pro	Leu	Leu	Ser	Asp	Pro	Ala	Leu	Asn	Leu
				185					190					195
Arg	Ile	Val	His	Leu	Val	Arg	Asp	Pro	Arg	Ala	Val	Leu	Arg	Ser
				200					205					210
Arg	Glu	Gln	Thr	Ala	Lys	Ala	Leu	Ala	Arg	Asp	Asn	Gly	Ile	Val
				215					220					225
Leu	Gly	Thr	Asn	Gly	Thr	Trp	Val	Glu	Ala	Asp	Pro	Gly	Leu	Arg
				230					235					240
Val	Val	Arg	Glu	Val	Cys	Arg	Ser	His	Val	Arg	Ile	Ala	Glu	Ala
				245					250					255
Ala	Thr	Leu	Lys	Pro	Pro	Pro	Phe	Leu	Arg	Gly	Arg	Tyr	Arg	Leu
				260					265					270
Val	Arg	Phe	Glu	Asp	Leu	Ala	Arg	Glu	Pro	Leu	Ala	Glu	Ile	Arg
				275					280					285
Ala	Leu	Tyr	Ala	Phe	Thr	Gly	Leu	Ser	Leu	Thr	Pro	Gln	Leu	Glu
				290					295					300
Ala	Trp	Ile	His	Asn	Ile	Thr	His	Gly	Ser	Gly	Pro	Gly	Ala	Arg
				305					310					315
Arg	Glu	Ala	Phe	Lys	Thr	Ser	Ser	Arg	Asn	Ala	Leu	Asn	Val	Ser
				320					325					330
Gln	Ala	Trp	Arg	His	Ala	Leu	Pro	Phe	Ala	Lys	Ile	Arg	Arg	Val
				335					340					345

Gln	Glu	Leu	Cys	Ala	Gly	Ala	Leu	Gln	Leu	Leu	Gly	Tyr	Arg	Pro	
				350					355					360	
Val	Tyr	Ser	Glu	Asp	Glu	Gln	Arg	Asn	Leu	Ala	Leu	Asp	Leu	Val	
				365					370					375	
Leu	Pro	Arg	Gly	Leu	Asn	Gly	Phe	Thr	Trp	Ala	Ser	Ser	Thr	Ala	
				380					385					390	
Ser	His	Pro	Arg	Asn											
				395											

&lt;210&gt; 6

&lt;211&gt; 504

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1272843CD1

&lt;400&gt; 6

Met	Glu	Ala	Thr	Gly	Thr	Trp	Ala	Leu	Leu	Leu	Ala	Leu	Ala	Leu	
1				5					10					15	
Leu	Leu	Leu	Leu	Thr	Leu	Ala	Leu	Ser	Gly	Thr	Arg	Ala	Arg	Gly	
				20					25					30	
His	Leu	Pro	Pro	Gly	Pro	Thr	Pro	Leu	Pro	Leu	Leu	Gly	Asn	Leu	
				35					40					45	
Leu	Gln	Leu	Arg	Pro	Gly	Ala	Leu	Tyr	Ser	Gly	Leu	Met	Arg	Leu	
				50					55					60	
Ser	Lys	Lys	Tyr	Gly	Pro	Val	Phe	Thr	Ile	Tyr	Leu	Gly	Pro	Trp	
				65					70					75	
Arg	Pro	Val	Val	Val	Leu	Val	Gly	Gln	Glu	Ala	Val	Arg	Glu	Ala	
				80					85					90	
Leu	Gly	Gly	Gln	Ala	Glu	Glu	Phe	Ser	Gly	Arg	Gly	Thr	Val	Ala	
				95					100					105	
Met	Leu	Glu	Gly	Thr	Phe	Asp	Gly	His	Gly	Val	Phe	Phe	Ser	Asn	
				110					115					120	
Gly	Glu	Arg	Trp	Arg	Gln	Leu	Arg	Lys	Phe	Thr	Met	Leu	Ala	Leu	
				125					130					135	
Arg	Asp	Leu	Gly	Met	Gly	Lys	Arg	Glu	Gly	Glu	Glu	Leu	Ile	Gln	
				140					145					150	
Ala	Glu	Ala	Arg	Cys	Leu	Val	Glu	Thr	Phe	Gln	Gly	Thr	Glu	Gly	
				155					160					165	
Arg	Pro	Phe	Asp	Pro	Ser	Leu	Leu	Leu	Ala	Gln	Ala	Thr	Ser	Asn	
				170					175					180	
Val	Val	Cys	Ser	Leu	Leu	Phe	Gly	Leu	Arg	Phe	Ser	Tyr	Glu	Asp	
				185					190					195	
Lys	Glu	Phe	Gln	Ala	Val	Val	Arg	Ala	Ala	Gly	Gly	Thr	Leu	Leu	
				200					205					210	
Gly	Val	Ser	Ser	Gln	Gly	Gly	Gln	Thr	Tyr	Glu	Met	Phe	Ser	Trp	
				215					220					225	
Phe	Leu	Arg	Pro	Leu	Pro	Gly	Pro	His	Lys	Gln	Leu	Leu	His	His	
				230					235					240	
Val	Ser	Thr	Leu	Ala	Ala	Phe	Thr	Val	Arg	Gln	Val	Gln	Gln	His	
				245					250					255	
Gln	Gly	Asn	Leu	Asp	Ala	Ser	Gly	Pro	Ala	Arg	Asp	Leu	Val	Asp	
				260					265					270	
Ala	Phe	Leu	Leu	Lys	Met	Ala	Gln	Glu	Glu	Gln	Asn	Pro	Gly	Thr	
				275					280					285	
Glu	Phe	Thr	Asn	Lys	Asn	Met	Leu	Met	Thr	Val	Ile	Tyr	Leu	Leu	
				290					295					300	
Phe	Ala	Gly	Thr	Met	Thr	Val	Ser	Thr	Thr	Val	Gly	Tyr	Thr	Leu	
				305					310					315	
Leu	Leu	Leu	Met	Lys	Tyr	Pro	His	Val	Gln	Lys	Trp	Val	Arg	Glu	
				320					325					330	



Glu	Leu	Asn	Arg	Glu	Leu	Gly	Ala	Gly	Gln	Ala	Pro	Ser	Leu	Gly	335	340	345
Asp	Arg	Thr	Arg	Leu	Pro	Tyr	Thr	Asp	Ala	Val	Leu	His	Glu	Ala	350	355	360
Gln	Arg	Leu	Leu	Ala	Leu	Val	Pro	Met	Gly	Ile	Pro	Arg	Thr	Leu	365	370	375
Met	Arg	Thr	Thr	Arg	Phe	Arg	Gly	Tyr	Thr	Leu	Pro	Gln	Gly	Thr	380	385	390
Glu	Val	Phe	Pro	Leu	Leu	Gly	Ser	Ile	Leu	His	Asp	Pro	Asn	Ile	395	400	405
Phe	Lys	His	Pro	Glu	Glu	Phe	Asn	Pro	Asp	Arg	Phe	Leu	Asp	Ala	410	415	420
Asp	Gly	Arg	Phe	Arg	Lys	His	Glu	Ala	Phe	Leu	Pro	Phe	Ser	Leu	425	430	435
Gly	Lys	Arg	Val	Cys	Leu	Gly	Glu	Gly	Leu	Ala	Lys	Ala	Glu	Leu	440	445	450
Phe	Leu	Phe	Phe	Thr	Thr	Ile	Leu	Gln	Ala	Phe	Ser	Leu	Glu	Ser	455	460	465
Pro	Cys	Pro	Pro	Asp	Thr	Leu	Ser	Leu	Lys	Pro	Thr	Val	Ser	Gly	470	475	480
Leu	Phe	Asn	Ile	Pro	Pro	Ala	Phe	Gln	Leu	Gln	Val	Arg	Pro	Thr	485	490	495
Asp	Leu	His	Ser	Thr	Thr	Gln	Thr	Arg							500		

&lt;210&gt; 7

&lt;211&gt; 229

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7472790CD1

&lt;400&gt; 7

Met	Asn	Ile	Arg	Asn	Ala	Gln	Pro	Asp	Asp	Leu	Met	Asn	Met	Gln	1	5	10	15
His	Cys	Asn	Leu	Leu	Cys	Leu	Pro	Glu	Asn	Tyr	Gln	Met	Lys	Tyr	20	25	30	35
Tyr	Leu	Tyr	His	Gly	Leu	Ser	Trp	Pro	Gln	Leu	Ser	Tyr	Ile	Ala	40	45	50	55
Glu	Asp	Glu	Asp	Gly	Lys	Ile	Val	Gly	Tyr	Val	Leu	Ala	Lys	Met	60	65	70	75
Glu	Glu	Glu	Pro	Asp	Asp	Val	Pro	His	Gly	His	Ile	Thr	Ser	Leu	80	85	90	95
Ala	Val	Lys	Arg	Ser	His	Arg	Arg	Leu	Gly	Leu	Ala	Gln	Lys	Leu	100	105	110	115
Met	Asp	Gln	Ala	Ser	Arg	Ala	Met	Ile	Glu	Asn	Phe	Asn	Ala	Lys	120	125	130	135
Tyr	Val	Ser	Leu	His	Val	Arg	Lys	Ser	Asn	Arg	Pro	Ala	Leu	His	140	145	150	155
Leu	Tyr	Ser	Asn	Thr	Leu	Asn	Phe	Gln	Ile	Ser	Glu	Val	Glu	Pro	160	165	170	175
Lys	Tyr	Tyr	Ala	Asp	Gly	Glu	Asp	Ala	Tyr	Ala	Met	Lys	Arg	Asp	180	185	190	195
Leu	Ser	Gln	Met	Ala	Asp	Glu	Leu	Arg	Arg	Gln	Met	Asp	Leu	Lys	200	205	210	

Glu Ser Val Glu Ser Thr Asn Val Gln Asp Ser Ser Glu Ser Ser  
 215 220 225  
 Asp Ser Thr Ser

<210> 8  
 <211> 347  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 7473944CD1

<400> 8  
 Met Lys Pro Ala Leu Leu Glu Val Met Arg Met Asn Arg Ile Cys  
 1 5 10 15  
 Arg Met Val Leu Ala Thr Cys Leu Gly Ser Phe Ile Leu Val Ile  
 20 25 30  
 Phe Tyr Phe Gln Ile Met Arg Arg Asn Pro Phe Gly Val Asp Ile  
 35 40 45  
 Cys Cys Arg Lys Gly Ser Arg Ser Pro Leu Gln Glu Leu Tyr Asn  
 50 55 60  
 Pro Ile Gln Leu Glu Leu Ser Asn Thr Ala Val Leu His Gln Met  
 65 70 75  
 Arg Arg Asp Gln Val Thr Asp Thr Cys Arg Ala Asn Ser Ala Thr  
 80 85 90  
 Ser Arg Lys Arg Arg Val Leu Thr Pro Asn Asp Leu Lys His Leu  
 95 100 105  
 Val Val Asp Glu Asp His Glu Leu Ile Tyr Cys Tyr Val Pro Lys  
 110 115 120  
 Val Ala Cys Thr Asn Trp Lys Arg Leu Met Met Val Leu Thr Gly  
 125 130 135  
 Arg Gly Lys Tyr Ser Asp Pro Met Glu Ile Pro Ala Asn Glu Ala  
 140 145 150  
 His Val Ser Ala Asn Leu Lys Thr Leu Asn Gln Tyr Ser Ile Pro  
 155 160 165  
 Glu Ile Asn His Arg Leu Lys Ser Tyr Met Lys Phe Leu Phe Val  
 170 175 180  
 Arg Glu Pro Phe Glu Arg Leu Val Ser Ala Tyr Arg Asn Lys Phe  
 185 190 195  
 Thr Gln Lys Tyr Asn Ile Ser Phe His Lys Arg Tyr Gly Thr Lys  
 200 205 210  
 Ile Ile Lys Arg Gln Arg Lys Asn Ala Thr Gln Glu Ala Leu Arg  
 215 220 225  
 Lys Gly Asp Asp Val Lys Phe Glu Glu Phe Val Ala Tyr Leu Ile  
 230 235 240  
 Asp Pro His Thr Gln Arg Glu Glu Pro Phe Asn Glu His Trp Gln  
 245 250 255  
 Thr Val Tyr Ser Leu Cys His Pro Cys His Ile His Tyr Asp Leu  
 260 265 270  
 Val Gly Lys Tyr Glu Thr Leu Glu Glu Asp Ser Asn Tyr Val Leu  
 275 280 285  
 Gln Leu Ala Gly Val Gly Ser Tyr Leu Lys Phe Pro Thr Tyr Ala  
 290 295 300  
 Lys Ser Thr Arg Thr Thr Asp Glu Met Thr Thr Glu Phe Phe Gln  
 305 310 315  
 Asn Ile Ser Ser Glu His Gln Thr Gln Leu Tyr Glu Val Tyr Lys  
 320 325 330  
 Leu Asp Phe Leu Met Phe Asn Tyr Ser Val Pro Ser Tyr Leu Lys  
 335 340 345  
 Leu Glu

<210> 9  
 <211> 218  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2244136CD1

<400> 9  
 Met Thr Pro Ala Pro Pro Pro Gly Ala Arg Pro Gly Ala Ala Ser  
 1 5 10 15  
 Leu Ala Gly Phe Ala Gly Val Ala Ser Leu Gly Pro Gly Asp Pro  
 20 25 30  
 Arg Arg Ala Ala Asp Pro Arg Pro Leu Pro Pro Ala Leu Cys Phe  
 35 40 45  
 Ala Val Ser Arg Ser Leu Leu Leu Thr Cys Leu Val Pro Ala Ala  
 50 55 60  
 Leu Leu Gly Leu Arg Tyr Tyr Tyr Ser Arg Lys Val Ile Arg Ala  
 65 70 75  
 Tyr Leu Glu Cys Ala Leu His Thr Asp Met Ala Asp Ile Glu Gln  
 80 85 90  
 Tyr Tyr Met Lys Pro Pro Gly Ser Cys Phe Trp Val Ala Val Leu  
 95 100 105  
 Asp Gly Asn Val Val Gly Ile Val Ala Arg Ala His Glu Glu  
 110 115 120  
 Asp Asn Thr Val Glu Leu Leu Arg Met Ser Val Asp Ser Arg Phe  
 125 130 135  
 Arg Gly Lys Gly Ile Ala Lys Ala Leu Gly Arg Lys Val Leu Glu  
 140 145 150  
 Phe Ala Val Val His Asn Tyr Ser Ala Val Val Leu Gly Thr Thr  
 155 160 165  
 Ala Val Lys Val Ala Ala His Lys Leu Tyr Glu Ser Leu Gly Phe  
 170 175 180  
 Arg His Met Gly Ala Ser Asp His Tyr Val Leu Pro Gly Met Thr  
 185 190 195  
 Leu Ser Leu Ala Glu Arg Leu Phe Phe Gln Val Arg Tyr His Arg  
 200 205 210  
 Tyr Arg Leu Gln Leu Arg Glu Glu  
 215

<210> 10  
 <211> 318  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 7474327CD1

<400> 10  
 Met Ala Ser His Ile Val Leu Asn Asn Gly Thr Lys Met Pro Ile  
 1 5 10 15  
 Leu Gly Leu Gly Thr Trp Asn Ser Pro Pro Gly Gln Val Thr Glu  
 20 25 30  
 Ala Val Lys Val Ala Ile Asn Val Gly Tyr Cys His Ile Asp Cys  
 35 40 45  
 Ala His Val Tyr Gln Asn Glu Asn Asp Val Gly Val Ala Ile Arg  
 50 55 60  
 Glu Lys Leu Arg Glu Gln Val Val Lys Cys Glu Glu Leu Phe Ile  
 65 70 75  
 Thr Ser Lys Leu Trp Cys Ala Tyr His Glu Lys Gly Leu Val Lys  
 80 85 90

```
<210> 11
<211> 1636
<212> DNA
<213> Homo sapiens
```

```
<220>  
<221> misc_feature  
<223> Incyte ID No: 2434655CB1
```

<400> 11						
gatcagtg	tgagggaact	gccatcatga	ggtctgacaa	gtcagctttg	gtattttctgc	60
tcctgcagct	cttctgtggt	ggctgtggat	tctgtgggaa	agtcctgggtg	tggccctgtg	120
acatgagcca	ttggcttaat	gtcaagggtca	ttctagaaga	gctcatagtg	agaggccatg	180
aggtaacagt	attgactcac	tcaaagcctt	cgtttaattga	ctacaggaag	ccttctgcat	240
tgaaatttga	ggtgggtccat	atgccacagg	acagaacaga	agaaaatgaa	atattttgtg	300
acctagctct	gaatgtcttg	ccaggcttat	caactgggca	atcagttata	aaattaaatg	360
atttttttgt	tgaataaaga	ggaactttaa	aaatgatgtg	tgagagcttt	atctacaatc	420
agacgcttat	gaagaagcta	caggaaacca	actacgatgt	aatgcttata	gaccctgtga	480
ttccctgtgg	agacctgatg	gctgagttgc	tgcagtcctc	ttttgtgctc	acacttagaa	540
tttctgtagg	aggcaatatg	gagcgaagct	gtgggaaact	tccagctcca	ctttcctatg	600
tacctgtgcc	tatgacagga	ctaacagaca	gaatgacctt	tctggaaaga	gtaaaaaatt	660
caatgctttc	attttgttgc	cacttttgga	ttcaggatta	cgactatcat	tttttgggaag	720
agttttatag	taaggcatta	ggaaggccca	ctacattatg	tgagactgtg	ggaaaagctg	780
agatatggct	aatacgaaca	tattgggatt	ttgaatttcc	tcaaccatac	caaccctaact	840
ttgagtttgt	tggaggattg	cactgtaaac	ctgccaaagc	tttgccctaag	gaaatggaaa	900
atttttgtcca	gagttcaggg	gaagatggta	ttgtgggtgtt	ttctctgggg	tcactgtttc	960
aaaatggttac	agaagaaaag	gctaatatca	ttgcttcacg	ccttgcccag	atcccacaga	1020
agggtgttatg	gaggtacaaa	ggaaaaaaac	catccacatt	aggagccaat	actcggctgt	1080
atgattggat	accccagaat	gatcttcttg	gtcatcccaa	aaccaaagct	tttatcactt	1140
atggtggaat	gaatgggatc	tatgaagcta	tttaccatgg	ggtccctatg	gtggggagttc	1200

```

ccatattttgg  tgatcagctt  gataacatag  ctcacatgaa  ggccaaagga  gcagctgtag  1260
aaataaactt  caaaactatg  acaagcgaag  atttactgag  ggctttgaga  acagtcatta  1320
ccgattcctc  ttataaagag  aatgctatga  gattatcaag  aattcaccat  gatcaacctg  1380
taaagcccct  agatcgagca  gtcttctgga  tcgagtttgt  catgcgccac  aaaggagcca  1440
agcacctgcy  atcagctgcc  catgacctca  cctgggtcca  gcactactct  atagatgtga  1500
ttgggttctt  gctgacctgt  gtggcaactg  ctatattctt  gttcacaaaa  tgttttttat  1560
tttctgtca  aaaatttaat  aaaactagaa  agatagaaaa  gaggggaatag  atctttccaa  1620
attcaagaaa  gacctg                                1636

```

```

<210> 12
<211> 2086
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 2516747CB1

```

```

<400> 12
cctacctctt  cctaggccca  cagccagtgc  ctttggagta  ctgaggcgcy  cacagagtcc  60
ttagcccgcy  gcaggcgcy  cagcccaggc  tgagatccgc  tgcttctgtg  gaagtgcaga  120
tggttgggca  gcgggtgctg  cttctagtgg  ccttctctct  ttctggggtc  ctgctctcag  180
aggctgccaa  aatcctgaca  atatctacac  tgggtggaag  ccattaccta  ctggttgacc  240
gggtgtctca  gattcttcaa  gagcatggtc  ataattgtac  tatgcttcac  cagagtggaa  300
agtttttgat  cccagatatt  aaagaggagg  aaaaatcata  ccaagttatc  aggtgggttt  360
cacctgaaga  tcatcaaaaa  agaattaaga  agcattttga  tagctacata  gaaacagcat  420
tggatggcag  aaaagaatct  gaagcccttg  taaagctaag  ggaaatattt  gggactcaat  480
gtagttatct  gctaagcaga  aaggatataa  tggattcctt  aaagaatgag  aactatgata  540
tggtatttgt  tgaagcattt  gatttctgtt  ctttctctgt  tgctgagaag  cttgtgaaac  600
catttgtggc  cattcttccc  accacattcg  gctctttgga  ttttgggcta  ccaagcccct  660
tgtcttatgt  tccagtattc  ccttctctgc  tgactgatca  catggacttc  tggggccgag  720
tgaagaattt  tctgatgttc  tttagtttct  ccaggagcca  atgggacatg  cagtctacat  780
ttgacaacac  catcaaggag  catttcccag  aaggctctag  gccagttttg  tctcatcttc  840
tactgaaagc  agagttgtgg  tttgttaact  ctgattttgc  ctttgatttt  gcccggcccc  900
tgcttcccaa  cactgtttat  attggaggct  tgatggaaaa  acctattaaa  ccagtaccac  960
aagacttggg  caacttcatt  gccaactttg  gggatgcagg  gtttgtcctt  gtggcctttg  1020
gctccatgtt  gaacacccat  cagtcccagg  aagtcctcaa  gaagatgcac  aatgcctttg  1080
cccacctccc  tcaaggagtg  atatggacat  gtcagagttc  tcattggccc  agagatgttc  1140
atttggccac  aaatgtgaaa  attgtggact  ggcttctctg  gagtgccttc  ctggctcacc  1200
ccagcatccg  tctttttgtc  actcatgggt  ggcagaacag  cgtaatggag  gccatccgtc  1260
atgggtgtgc  catgggtggg  ttaccagtca  atggagacca  gcatggaaac  atggctccag  1320
tagtagccaa  aaattatggg  gtctctatcc  ggttgaatca  ggtcacagcc  gacacactga  1380
cattacaatt  gaaacaagtc  atagaagaca  agaggtacaa  gtcggcagtg  gtggcagcca  1440
gtgtcatact  gcactctcag  cccctgagcc  ccgcacagcg  gctgggtggc  tggatcgacc  1500
acatcctcca  gactggggga  ggcagcgacc  tcaagcccta  tgccctccag  cagccttggc  1560
atgagcagta  cctcatgtat  gtcttttgtt  ttctgtctgg  gctcactctg  ggcactatgt  1620
ggctttgtgg  gaagctgctg  ggtgtgggtg  ccaggtggct  gcgtggggcc  aggaaggtga  1680
agaagacatg  aggctagggt  tagccttggg  tgaggggagg  gcatccctgg  tcctttgaag  1740
gttctcccca  ccccgacaca  cgccacccct  ctgttctctc  ttcagctcca  cctgccactg  1800
atcctgcaac  ttgcttcttt  ctattctctg  cctctgttta  gaaatcttca  cacaccactg  1860
aggcttcttg  acttgccttc  tgtgacttga  attcccagct  cagatacaaa  ttttcacctg  1920
ccagccctgc  ctctctcttt  ctcccttttc  ctagacacag  gactctgaca  acttcatcct  1980
ccttggttag  atgacttccc  agtttccagt  cccattttct  cttctatca  cttttcataa  2040
aaaaactcag  gaaatatttg  acatatcttc  catttcaaat  tcttcc                                2086

```

```

<210> 13
<211> 1814
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 7472775CB1

```

&lt;400&gt; 13

```
cttggacgag ggggagagtg gagaagagaa gagaacaaaa agtaacttca ggatggggac 60
cagtgaagta cttgcccgtt gtacccagaa tcaaacccca agtttcacat gcctgaggat 120
gtacgagaaa aaaaggaaaa tcttctactc aattctgaga gatctactag gctcttaaca 180
aagaccagtc attcacaagg aggggatcaa gctttaagta agtccacagg gtcaccaaca 240
gagaagttga ttgaaaaacg tcaaggagct aagactgttt ttaacaagtt cagcaacatg 300
aattggccag tggacattca ccctttaaac aaaagtttag tcaaagataa taaatggaag 360
aaaactgagg agaccaaga gaaacgaagg tctttccttc aggagttttg caagaaatac 420
ggtggggtga gtcacatca gtcacatctt ttccatacag tatccagaat ctatgtagaa 480
gataaacaca aaatcttata ttgtgaggta cctaaggctg gctgttccaa ttggaaaaga 540
attctgattg tactaaatgg attggcttcc tctgcataca acatctccca caatgctgtc 600
cactacggga agcatttgaa gaagctagat agctttgacc taaaagggat atataccgcg 660
ttaataactt acaccaaagc tgtgtttgtt cgtgatccca tggaaagatt agtatcagcc 720
tttagggaca aatttgaaca ccccaatagt tattaccatc cagtattcgg aaaggcaatt 780
atcaagaaat atcgaccaa tgcctgtgaa gaagcattaa ttaatggatc tggagtcaag 840
ttcaaagagt ttatccacta cttgctggat tcccaccgtc cagtaggaat ggacattcac 900
tgggaaaagg tcagcaact ctgctatccg tgtttgatca actatgattt tgtagggaaa 960
tttgagactt tggagaaga tgccaattac tttttacaga tgatcgggtc tccaaaggag 1020
ctgaaatttc ccaactttaa ggataggcac tcttccgatg aaagaaccaa tgctcaagtc 1080
gtgagacagt atttaaagga tctgactaga actgagagac aattaatcta tgacttttat 1140
tacttggact atttaatgtt taattatata actccatttt tgtagtttgc attcattttc 1200
taaaaccctg tatatactta atgatgataa gttcaaatac gctgtaattt ttctataatt 1260
ctctgtatga cagaaattta accaagtga gttgtcttga tttaatgtag atttttacca 1320
aatagtatga caccaattgg cacaagttta taggaaaatc acctacagga gatgtaaaaca 1380
acttgagttg ctctaaaatg tttggaaaag agctgctttt gcattatgaa ttatattgtt 1440
gaagcaataa cctagccagc tgttgcatca gctaaagcag cctcttgcaa tggtaggaaa 1500
aaaggatctc aaatagcatg agtgtatgtc tatatcctga aattttattgt ctaaaatgca 1560
tgaatatatt tttagcagtc tgtggcatat taatcaaact gttgaattgt tttcttacac 1620
cctggaaatc tttctatcaa ctataatgat aaatccattt tgaagtgata ttttggactt 1680
aggcatttta ctttagattg gaaggcatca tgtgatttac aatatgagaa tatagcagaa 1740
aaaccagatg aggctgtggc tttttatatt caacagccaa taaaaaatgc acaacatgct 1800
aagatcaaag caaa 1814
```

&lt;210&gt; 14

&lt;211&gt; 1650

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7473323CB1

&lt;400&gt; 14

```
atggccattg atgcactggt ctctctgtgt cttcctgagg tcatcagaat aaagttcaac 60
atcaggccaa ggcagcccca ccacgacctc ccaccaggcg gctcccagga tgggtgacttg 120
aaggaaacca cagagagggt cactcgggac ttatccagtg gggccccgag gggccgcaac 180
ctgccagcgc ctgaccagcc tcaacccccg ctgcagaggg gaaccgctct gcggtccgcg 240
cagcgcgcgtc gccgtctgct catcaagaaa atgcagctg cggcgaccat cccggccaac 300
agctcggagc cgcccttcat ccggccggga ccggggacgc tggatggccg ctgggtcagc 360
ctgcaccgga gccagcagga gcgcaagcgg gtgatgcagg aggcctgcgc caagtaccgg 420
gcgagcagca gccgcggggc cgtcacgccc cgccacgtgt cccgtatctt cgtggaggac 480
cgccaccgcg tgctctactg cgaggtgccc aaggccggct gctccaattg gaagcgggtg 540
ctcatggtgc tggccggcct ggccctcgtcc actgcgaca tccagcaca caccgtccac 600
tatggcagcg ctctcaagcg cctggacacc ttgcagcgcc aggggtatctt gcaccgtctc 660
agcacctaca ccaagatgct ctttgtccgc gagcccttcg agaggctggt gtccgccttc 720
cgcgacaagt ttgagcacc caacagctac tatcaccgg tcttcggcaa ggccatcctg 780
gccccgtacc gcgccaatgc ctctcgggag gccctgcgga cgggctctgg ggtgcgtttt 840
cccagattcg tccagtacct gctggacgtg caccggcccc tggggatgga cattcactgg 900
gaccatgtca cgcggctctg cagccctctc ctcatcgact acgatttcgt aggcaagttc 960
gagagcatgg aggagcatgc caacttcttc ctgagctca tccgcgcgcc gcggaacctc 1020
accttcccc gggtcaagga ccggcactcg caggaggcgc ggaccacagc gaggaatcgc 1080
caccagtact tcgccaact ctcgccctcg caaaggcagc gcacctacga cttctactac 1140
atggattacc tgatgttcaa ctattccaag ccctttgcag atctgtactg aggggcgcgc 1200
```

cagctggccg	gggcccgcct	gccccgggtca	ctcacctgtg	ctccccgggca	tcctcctgtc	1260
cctggctcct	catectggga	gcaacagggc	tctgaggacg	tgaggagcca	tcgctgtggg	1320
aggcagcagg	ccccgggtgg	ggggcagagg	cgcccagcct	tggatgggga	ccccagcccc	1380
tggcctgtac	ctgtttcctc	attccttggc	tgaggagagag	gctgagaact	gggcagacac	1440
ccctggagct	cagccgacag	ttttgatgag	caggggaagtc	tgaggcccag	aggacggggg	1500
gcccagcggg	aagggatgtc	ccgcactccc	ttagccattg	ccttggacca	aaccacgtgg	1560
tttgacgctt	ttctacgagc	caggggggag	gttcccttgg	attaaggttc	caaataaagc	1620
acatggtttc	cagaacaaaa	aaaaacaaaa				1650

&lt;210&gt; 15

&lt;211&gt; 1647

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7472777CB1

&lt;400&gt; 15

atgtggctgc	cgcgctctc	cagcacagca	gtgaccgcgc	tcctcctggc	gcagaccttc	60
ctcctcctct	ttctggtttc	ccggccaggg	ccctcgtccc	cagcaggcgg	cgaggcgcg	120
gtgcatgtgc	tgggtgctgc	ctcgtggcgc	tcgggctcgt	ccttcgtggg	ccaactcttc	180
aaccagcacc	ccgacgtctt	ctacctaatg	gagcccgctg	ggcacgtgtg	gaccaccctg	240
tcgcagggca	gcgccgaac	gctgcacatg	gctgtgcgcg	acctggtgcg	ctccgtcttc	300
ctgtgcgaca	tggacgtgtt	tgatgcctat	ctgccttggc	gccgcaacct	gtccgacctc	360
ttccagtggg	ccgtgagccg	tgactgtgc	tcgccaccgc	cctgcagtgc	ctttccccga	420
ggcgccatca	gcagcgaggc	cgtgtgcaag	ccactgtgcg	cgcggcagtc	cttcaccctg	480
gcccgggagg	cctgccgctc	ctacagccac	gtggtgctca	aggaggtgcg	cttcttcaac	540
ctgcaggtgc	tctacccgct	gctcagcgac	cccgcgctca	acctacgcat	cgtgcacctg	600
gtgcgcgacc	cgcgggccgt	gctgcgctcc	cgggagcaga	cagccaaggc	tctggcgctg	660
gacaacggca	tcgtgctggg	caccaacggc	acgtgggtgg	aggccgacct	cgccctgcgc	720
gtggtgcgcg	aggtgtgccc	tagccacgta	cgcctcgccg	aggccgccac	actcaagccg	780
ccaccctttc	tgcgcggccg	ctaccgcctg	gtgcgcttcg	aggacctggc	gcgggagccg	840
ctggcagaaa	tcctgtgcgt	ctacgccttc	actgggctca	gtctcacgcc	acagctcgag	900
gcctggatcc	ataacatcac	ccacggatct	ggacctggtg	cgcgccgcga	agccttcaag	960
acttcgtcca	ggaatgcgct	caacgtctcc	caggcctggc	gccatgcgct	gccctttgcc	1020
aagatccgcc	cggtgcagga	actgtgcgct	ggtgcgctgc	agctgctggg	ctaccggcct	1080
gtgtactctg	aggacgagca	gcgcaacctc	gcccttgatc	tgggtgctgc	acgaggcctg	1140
aacggcttca	cttgggcata	atccaccgcc	tcgcaccccc	gaaattagtg	gaggccacag	1200
ttgtagcagg	cgctaggccc	gggaggagag	tgcatggtgc	agagggggct	ggggcgacag	1260
gagaagcagg	tcctatatatt	gaccaaggag	tttgtgagaa	cctgcgtgct	gtccctttgc	1320
ttcggacctc	cgctcttgcc	cgggagaaaag	cccaggccag	cctgctggac	aagcagagac	1380
catgagaagg	agagttcagg	ggtcccaaac	caggccatcc	tagaccagcc	agctccagct	1440
gatccgcacg	cagccacttc	ggctaccttc	tactggccaa	agggagtccc	agggctcacc	1500
cagattcaga	ggtggggaaa	ctgagtccac	cacttgagaa	gagtagctat	aaagacatat	1560
gagcgaggcc	agctgagccc	agcactgcgg	ccaagtcgaa	gctttaggag	caataaaaagt	1620
gcttattgtg	tttcagtcaa	aaaaaaa				1647

&lt;210&gt; 16

&lt;211&gt; 2620

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1272843CB1

&lt;400&gt; 16

cgcgagagacc	tgggagagga	gaaggagccg	acctgccgag	atggaggcga	ccggcacctg	60
ggcgctgctg	ctggcgctgg	cgctgctcct	gctgctgacg	ctggcgctgt	ccgggaccag	120
ggcccaggc	cacctgcccc	ccgggcccac	gccgctacca	ctgctgggaa	acctcctgca	180
gctacggccc	ggggcgctgt	attcagggct	catgcgctg	agtaagaagt	acggaccggg	240
gttcaccatc	tacctgggac	cctggcgggc	tgtggtggtc	ctgggtgggc	aggaggctgt	300

```

gcgaggagcc ctgggaggtc aggctgagga gttcagcgcc cggggaaccg tagcgatgct 360
ggaagggact tttgatggcc atgggggtttt cttctccaac ggggagcggt ggaggcagct 420
gaggaagtgt accatgcttg ctctgcgggg cctgggcatg gggaagcgag aaggcgagga 480
gctgatccag gcggaggccc ggtgtctggt ggagacattc caggggacag aaggacgccc 540
attcgatccc tccctgctgc tggcccagge cacctccaac gtagtctgct ccctcctctt 600
tggcctccgc ttctcctatg aggataagga gttccaggcc gtgggtccggg cagctggtgg 660
taccctgctg ggagtcagct cccaggggggg tcagacctac gagatgttct cctggttcct 720
gcggccccctg ccaggccccc acaagcagct cctccaccac gtcagcacct tggctgcctt 780
cacagtccgg caggtgcagc agcaccaggg gaacctggat gcttcggggc ccgcacgtga 840
ccttgctgat gccttcctgc tgaagatggc acaggaggaa caaaacccag gcacagaatt 900
caccaacaag aacatgctga tgacagtcat ttatttgctg ttgctggga cgtgacgggt 960
cagcacaccg gtcgggtata ccctcctgct cctgatgaaa taccctcatg tccaaaagt 1020
ggtacgtgag gagctgaatc gggagctggg ggctggccag gcaccaagcc taggggaccg 1080
taccgcctc ccttacaccg acgcggttct gcatgaggcg cagcggtgc tggcgctggt 1140
gcccattggg ataccccgca ccctcatgag gaccaccgc ttccgagggt acaccctgcc 1200
ccagggcacg gaggtcttcc ccctccttgg ctccatcctg catgacccca acatcttcaa 1260
gcaccagaa gagttcaacc cagaccgttt cctggatgca gatggacggg tcaggaagca 1320
tgaggcgctc ctgccccttc ccttagggaa gcgtgtctgc cttggagagg ccctggcaaa 1380
agcggagctc ttccctcttct tcaccacat cctacaagcc ttctccctgg agagccctg 1440
cccgcggac accctgagcc tcaagcccac cgtcagtggc cttttcaaca ttccccagc 1500
cttcagctg caagtccgtc ccactgacct tcactccacc acgcagacca gatgaaggaa 1560
ggcaacttgg aagtgggtgg tgcccaggac ggtgcctcca gcctcaacag tgggcatgga 1620
cagggttaat gtctccagag tgtacactgc aggagccac atttacacgc ctgcagttgt 1680
tttcggaggt ctgtcccacg gccacacgc tcaactgact catgctgcta agatgcacaa 1740
ccgcacacc atacacaact acaagggcca caaagcaact gctgggttag ctttcacag 1800
acataaatat agtccatctg caatcacaag cacatagcca ggtaaccac caactcccc 1860
ggatctgcag cccacacgtg ggagtctggc tgtcacctc acaagccaca gaaacggcca 1920
cacatgttca cagctcacac gccctctcca ttcatcgaac ttctcagtg 1980
ggtgcctggc acagggaaca gcatgcccc tccgggtca tgccaccag agactgtcgc 2040
tgtctatggc cccaactcat gctccctctc ttgggtacac cactctcca gcctgtgacc 2100
accgatgtcc acacaccccc aaccacttgt ccacacagct acccagtac gacatcgctc 2160
tggctccca gagtatcttc ccactgagac acgcgcgcc cagagaggca cagtccccag 2220
ccacctctgc aactgcagcc ctcaatrac cctttttaag caccctgatt ctaccaaagt 2280
caaacacatc tgggtctgag attatgcaca gagactttgg acatacgagg accctcagac 2340
cggaggaaca cctgcccac ccaaacagc gcttatgtaa ccacgtggaa agcggcccc 2400
gctgccccct cacacacaca tacacacta ctgatctaca gccctgttc ggcgtcagag 2460
tccccactag acccagtga aggggttag gaccaagtag gggccagttt ccaattcacc 2520
ctgtcaggga gtgagccgga tctgacgttc cttgtgactt aagggtccgg cttgggaatt 2580
aaagtttgtt tctggccttt agcctaaaaa aaaaaaaaaa 2620

```

&lt;210&gt; 17

&lt;211&gt; 690

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7472790CB1

&lt;400&gt; 17

```

atgaacatcc gcaacgctca gccagacgac ctgatgaata tgcaacactg caacctcctt 60
tgcttcctc agaactacca gatgaaatac tatttatatc atggccttct ctggccccag 120
ctttcttaca tcgctgagga tgaggacggg aagattgtgg gctatgttct ggccaaaagt 180
gaggaggaac cagatgatgt cccgcattgc catatcacct cactggccgt gaagcggtta 240
caccggcgcc tcggcctggc ccagaagctg atggaccagg cctccagggc catgatagag 300
aactttaacg ccaaatacgt gtctctgcac gtcaggaaga gtaaccggcc agccttgac 360
ctttattcta acacctcaa ctttcagatt agtgaggtgg aacctaaata ctatgcagat 420
ggggaagatg cttatgctat gaagcgggat ctctcgagga tggcagatga gctgagacga 480
caaatggacc tgaagaaggg cgggtatgtg gtccctgggt ccaggagaa ccaggagacc 540
cagggcagca cactttctga ttctgaagag gcctgtcagc aaaagaaccc ggctaccgaa 600
gaaagtggca gtcagacgaa agaacctaa gagtctgtgg agagcaccaa cgtccaggac 660
agctcagaaa gctcggattc cacctcctag 690

```



<210> 18  
<211> 1510  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 7473944CB1

<400> 18  
gcgggccgcga agcgactccg atcctccctc tgagccttgc tcagctctgc cccgcgcctc 60  
ccggggtccg gtccgcgcgg cgggggtccct gctcctgcgc cccggggcgcg cttcccggac 120  
atcccgggtcc ccgcagccag gacaaagcca tgaagccagc gctgctggaa gtgatgagga 180  
tgaacagaat ctgccggatg gtgctggcca cttgcttggg atcctttatc ctgggtcatct 240  
tctattttcca aatcatgcgg aggaatccct ttgggtgtgga catctgctgc cggaagggggt 300  
cccgaagccc cctgcaggaa ctctacaacc caatccagct ggagctctca aacactgctg 360  
tcctgcacca gatgcggcgg gaccaggtga cagacacgtg ccgagccaac agcgccacaa 420  
gccgtaagcg gagggtgctg accccaacg acctgaagca cttgggtggg gatgaggacc 480  
acgagctcat ctactgctac gtgcccgaagg tggcctgcac caactggaag cgggtcatga 540  
tggtcctgac cgggcggggg aagtacagcg accccatgga gatcccggcc aacgaggcac 600  
acgtctccgc caacctgaag accctgaacc agtacagcat ccagaaaatc aaccaccgct 660  
tgaaaagcta catgaagttc ctgtttgtcc gggagccctt cgagaggcta gtgtccgcct 720  
accgcaacaa gttcaccag aagtacaaca tctccttcca caagcggtag ggcaccaaga 780  
tcatcaaacy ccagcgggaag aacgccaccc aggaggccct gcgcaaaggg gacgatgtca 840  
aattcgagga gtttgtggcc tatctcatcg acccacacac ccagcgggag gagcctttca 900  
acgaacactg gcaaacgcgc tactcactct gccatccctg ccacatccac tatgacctcg 960  
tgggcaagta cgagacactg gaagaggatt ctaattacgt cctgcagctg gcaggagtgg 1020  
gcagctacct gaagtcccc acctatgcaa agtctacgag aactactgat gaaatgacca 1080  
cagaattctt ccagaacatc agctcagagc accaaacgca gctgtacgaa gtctacaaac 1140  
tcgatttttt aatgttcaat tactcagtg ccaagctacct gaaattggaa taaaggggggt 1200  
ggggagaggg agagaatcat gctttttaat ttaagatttt tatttgtcaa agaattata 1260  
tggaatttgg gttattttgt aaattaatat ttctttgggg atgatgctgc gagcagcata 1320  
gtgagaatta tttaaaatcc ttcgtaggga aggacagctg tctttgcagg ggaatatagga 1380  
tgggtcgtcc ttgtctgtag aagtgaatac tgcaacactg tctcaaagggt ttcttgtgtt 1440  
ctggtgaatt ccatgaattg tgcattccat aaattctaata taatattatt tatagttatt 1500  
taaaaaaaaaa 1510

<210> 19  
<211> 3701  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2244136CB1

<400> 19  
gagctcgggt cactatgtac gtgcgcagtg tgctggcaag gggagaagta gaaagcacc 60  
tgagatgacc ccgggtcctc caccaggagc gcggccgggc ggggctccc tagcgggctt 120  
cgccgggggtg gcgtctctgg ggcctgggga ccccgcgcgc gccgctgacc cgcgccctct 180  
gccccagcg ctgtgcttcg ccgtgagccg ctcgctgctg ctgacgtgcc tgggtgccggc 240  
cgcgctgctg ggcctgcgct actactacag ccgcaagggt atccgcgcct acctggagt 300  
cgcgctgcac acggacatgg cggacatcga gcagtactac atgaagccgc ccggctcctg 360  
cttctgggtg gccgtgctgg atggcaacgt ggtgggcatt gtggctgcac gggcccacga 420  
ggaggacaac acggtggagc tgctgcggtg gtctgtggac tcacgtttcc gaggcaagg 480  
catcgccaag gcgctgggcc ggaagggtgt ggagttcgcc gtgggtgcaca actactccgc 540  
ggtggtgctg ggcacgacgg ccgtcaagggt ggcgcgccac aagctctacg agtcgctggg 600  
cttcagacac atgggcgcca gtgaccacta cgtgctgccc ggcattgacc tctcgctggc 660  
tgagcgcctc ttcttcagg tccgtacca ccgctaccgc ctgcagctgc gcgaggagt 720  
accgcccgcg ctgcgccgcc cgcgccccc ggcgcctgt ccgcctttgc ccgcctgccc 780  
gccgcccgcg gcggcctgct ttcagacgct cacttggcgt ttgtgttggg tttccccctt 840  
tcaacatcct gcggtgtct ggctggttcc ggggggtgcg gggctgttgt tcttcggcga 900  
cactttgggtg ggggtgggtt gttgtgcgcc atagcccctc gcccttcccc acctgcctgg 960

```

gcggtcttggc acctgaagag tggcatcttg gaccaccgag gctgtccatg acgctgccc 1020
gcccggccgac actggggaag gcccagcctt gctcaccaag cacagaacct ctgcagcaga 1080
tcccgggggac aggtctccggc cccgcctgag gccccagcgc cactgcctgt ggaggccga 1140
gctggccacg gcgtctgttt gctccgcgca tgccgagggg gtggccccggc tgagcatgcc 1200
gcatacacac agccccgccc tgccgccttg cccagacttg acccgagac ccgggctggt 1260
gagcgccctt gtccccagcc cccagctggc tgtgggaggg cctgccccctg cccccacct 1320
ctggaggggc tggctctgcc cgcgcgcgcc ggctctgtcc acacctgctt tgctctgacg 1380
ccctccattt ctctggctcc ggccccctcc ctgctgggc tgtgctgact ggtgtcatca 1440
cccagggtgac tcccatggcg tccgtggcac agccagggtg ggggtccatg ggaccctct 1500
ccccagtgcc cactggatcg tctggcctc tcccagatgt ccccggggac ctctgcctc 1560
tggctgacgg tccacctgt gaatcttatc agcccaggct gctgccaaca gcgcccagcc 1620
cacagcttct cccagcctga aaccaacaca ttttctaata agttatttag acagaatagg 1680
actctgcatg actttaattc ttgggacaaa acggtagttt gtaccctaag acacagtttc 1740
tggccagtg tgatgggggt ggggtggcgg gtgggtggag cgttttgctg ttggaaacct 1800
ggagggaac cctgattgga tgcatttcc tgccatggag cagcctccc agcctggcc 1860
tgcagggtgg gcagggtgg gggcaaggga gtccgcagcc tccgggagga ggggcagggc 1920
gctgccttgg gctgggtggg aagaggggtg gccgcctcg cttccgctgg ccatgctct 1980
ggtctctcct tctgaggtc acaggcaggc gctgccttg acggggggcg ggggggtgg 2040
cctggaaggg gagacagagg tggagggtgg cacaggctgc acattcagct tagaagtgg 2100
cctggctttg gtggcaggag aagaataaac acttgcccag accctttgt gtgggggaat 2160
tggggagggg tctggcagg cagggtgggc cacggaactg ggtcccaggc atcaaggcca 2220
cgtgcagggc catggaggga tcttctcac gaggcgcttc agaagcgagc gaagggacag 2280
agaagccctg cgtccaagg ccttttgcct tgtagcaat tgagggtgtg agagcactgt 2340
acagacccca ctcccctgta cttctctccc tggaggtgcc cgggtcccgc ttggggatgg 2400
gagttttgta gactgtacag aaatcgac cctattttct tgcagctcag attttgtta 2460
tctggaatat acagacagac gtaaaagtgt ttagcaaaat ggaaacaaac agttgtgct 2520
tttctctctt ttggtttggt ttgggtttgg cctggggctg gtcccagttg gtgcggggca 2580
tgctgggggc aggaggggca gggcgggcca ggtggagtca ggtcttgggg gtgtcatgtc 2640
gggggtgctg cagcgtccct tggctcctgt cctttggagc ctggggtcc tgggggtgcag 2700
ggtgcttggg ggtggctgtt ggagcccacc gacgccaggg cagggcctgg agggccaggg 2760
actgccagtg tctccttgat attgatccta gcagatcccc ctctggggg ttctgagagt 2820
cctgggcagt gtggccttct ctcatctctg tggcatctgc gccctgagt gacctcttcc 2880
ttggctgcac tgcctgtgg gtggtgagac gcttggcctt tttgttgc gccaggactt 2940
ggtagagatg gcagggaagg tgtgcggggg ggttgtgtgc agaaccctgc cggccctgag 3000
gtgagcagag gactgggtgt gcctgccagg ctggggcgga gctgcccgga acccttgcca 3060
ggccaatgtg tagcttgggt gcgcccttga aggccactgg ggggtagggt tgcctcccc 3120
cggggctgga ggccgggtgc ctggtgggtg ggcctgacct gcccacctc atccctccag 3180
cctgggatct cactgctctg cacctgtgca ctgacgtgaa tttatacgt ttgtagaaac 3240
tctgatgtaa ctcttctac ctctgaagcg cctcctggg ccctgctgtc acgtggctgg 3300
tgcggtctcc ccgagtgtg gccccctgcc cgctccccat gggcacccac cactcaaaact 3360
ctgcgtgggt agggccgggg aaagccaggc cgggccctag cactgctcag gggcctgggg 3420
ctcctgggat ttctgtgtgt ttggaagcct ctgttttgg agtggggggc gtggaaggcg 3480
ggaggggctg acagtgcctg gggccagagc tggggcgaag aggcacacct cactggagca 3540
gcactgcagg cctgacgaga ggctcccggc agctgagggc ttaagcgct cgtgatgtcc 3600
ccaatggccc aagccccctg tctgcttgc tgggggtggg ttgccccgc gattgcaactg 3660
tcccctgggt ttcttcgacc agctcctccg gggagggcct c 3701

```

&lt;210&gt; 20

&lt;211&gt; 960

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 7474327CB1

&lt;400&gt; 20

```

gcagccatgg ccagccacat tgtgtcaca aatggcacca agatgcccac cctggggcta 60
ggcacctgga attccccctc agggcaggta actgaggctg tgaagggtgg cattaatgtt 120
gggtactgcc acatcgactg tgcccaggtg taccagaatg agaatgacgt ggggggtggc 180
attcgggaga agctcaggga gcagggtgag aagtgtgagg agctcttcat caccagcaag 240
ctgtggtgag cgtaccatga gaagggcctg gtgaaaggag cctgccagaa gatgctcatt 300
gacctgaagc tggactacct ggacctctac cttattcgct ggccaaccag cttcaagcct 360

```

gggaaggaat	ttttccatt	ggatgagcca	ggtaatggta	atgtggttcc	cagtaacagt	420
aacatttctgg	acacatgggc	gggcatggaa	gagctggtgg	atgaagggt	ggtgaaagct	480
attggcatct	ccaacttcaa	ccatctccag	gttgagagga	tcttaaacia	acctgactta	540
aagtataagc	cggtggttaa	tcagattgag	tgccaccctg	acctcactca	ggagaagtta	600
atccagtact	gccagtccaa	aggcatcatg	gtgactgctt	acagctcctt	cagctcccc	660
gacaggccct	gggccaagcc	tgaggaccct	tccctcctgg	aagatcccag	gatcaaggcg	720
atcacagcca	aacacaataa	aactacagcc	caggttctga	tctggttccc	catgcagagg	780
aacttggtgg	tgatcccaa	gtctgtgaca	ccagaatgca	ttgctgagaa	ctttaaggtc	840
ttcaactttg	aactgaacag	ccaggatatg	accaccttat	tcagttacia	cagaaactgg	900
agggctctgtg	ccttggtgag	ctgtgcctcc	cacaaggatt	accccttcca	tgaagagttt	960



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number  
**WO 01/079468 A3**

(51) International Patent Classification<sup>7</sup>: C12N 15/52,  
9/00, 15/63, 5/10, A01K 67/027, C12P 21/00, C07K  
16/40, C12Q 1/68, A61K 38/43, C12Q 1/00, G01N 33/573

(21) International Application Number: PCT/US01/11869

(22) International Filing Date: 12 April 2001 (12.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/197,590	13 April 2000 (13.04.2000)	US
60/198,403	19 April 2000 (19.04.2000)	US
60/200,185	28 April 2000 (28.04.2000)	US
60/202,234	5 May 2000 (05.05.2000)	US
60/203,509	11 May 2000 (11.05.2000)	US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US	60/198,403 (CIP)
Filed on	19 April 2000 (19.04.2000)
US	60/197,590 (CIP)
Filed on	13 April 2000 (13.04.2000)
US	60/200,185 (CIP)
Filed on	28 April 2000 (28.04.2000)
US	60/202,234 (CIP)
Filed on	5 May 2000 (05.05.2000)
US	60/203,509 (CIP)
Filed on	11 May 2000 (11.05.2000)

(71) Applicant (for all designated States except US): INCYTE GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): POLICKY, Jennifer, L. [US/US]; 1511 Jarvis Court, San Jose, CA 95118 (US). HAFALIA, April [US/US]; 2227 Calle de Primavera, Santa Clara, CA 95054 (US). BURFORD, Neil [GB/US]; 105 Wildwood Circle, Durham, CT 06422 (US). RING, Huijun, Z. [US/US]; 625 Orange Avenue, Los Altos, CA 94022 (US). LAL, Preeti [IN/US]; P.O. Box 5142, Santa Clara, CA 95056 (US). TRIBOULEY, Catherine, M. [FR/US]; 1121 Tennessee Street, #5, San Francisco, CA 94107 (US). YAO, Monique, G. [US/US]; 111 Frederick Court, Mountain View, CA 94043 (US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA

94087 (US). TANG, Y., Tom [US/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). PATTERSON, Chandra [US/US]; 490 Sherwood Way #1, Menlo Park, CA 94025 (US). DAS, Debopriya [IN/US]; Apartment 3, 1179 Bonita Avenue, Mountain View, CA 94040 (US). SANJANWALA, Madhu, S. [US/US]; 210 Sylvia Court, Los Altos, CA 94024 (US). GANDHI, Ameena, R. [US/US]; 837 Roble Avenue, #1, Menlo Park, CA 94025 (US). REDDY, Roopa [US/US]; 1233 W. McKinley Avenue, #3, Sunnyvale, CA 94086 (US). KHAN, Farrah, A. [IN/US]; 333 Escuela Avenue, #221, Mountain View, CA 94040 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). RAMKUMAR, Jaya [IN/US]; 34359 Maybird Circle, Fremont, CA 94555 (US). GRIFFIN, Jennifer, A. [US/US]; 181 Irene Court, #11, Belmont, CA 94002 (US). AU-YOUNG, Janice [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US).

(74) Agents: HAMLET-COX, Diana et al.; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:  
18 July 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DRUG METABOLIZING ENZYMES

(57) Abstract: The invention provides human drug metabolizing enzymes (DME) and polynucleotides which identify and encode DME. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with aberrant expression of DME.

WO 01/079468 A3

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/11869

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/52 C12N9/00 C12N15/63 C12N5/10 A01K67/027  
C12P21/00 C07K16/40 C12Q1/68 A61K38/43 C12Q1/00  
G01N33/573

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

SEQUENCE SEARCH, EPO-Internal, WPI Data, PAJ, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EM_HUM [Online] EMBL; 16 January 2000 (2000-01-16) WATERSTON R.H.: "Homo sapiens chromosome 4 clone RP11-468N14, complete sequence" retrieved from EBI, accession no. AC021146 Database accession no. AC021146 XP002186102 the whole document	3,12
A	WO 97 44466 A (ENDORECHERCHE INC ;LEVESQUE ERIC (CA); BEAULIEU MARTIN (CA); BELAN) 27 November 1997 (1997-11-27) Seq Id Nos 1,2 abstract; claims page 1, line 20 -page 2, line 4 --- -/--	1-45,55

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

8 January 2002

Date of mailing of the international search report

07.05.02

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

CEDER O.

# INT. NATIONAL SEARCH REPORT

International Application No

PCT/US 01/11869

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 905 232 A (KIRIN BREWERY) 31 March 1999 (1999-03-31) abstract page 1, line 1 -page 5, line 24 ---	
A	EP 0 463 395 A (HOFFMANN LA ROCHE) 2 January 1992 (1992-01-02) ---	
A	KIMURA S ET AL: "THE HUMAN DEBRISOQUINE 4-HYDROXYLASE (CYP2D) LOCUS: SEQUENCE AND IDENTIFICATION OF THE POLYMORPHIC CYP2D6 GENE, A RELATED GENE, AND A PSEUDOGENE" AMERICAN JOURNAL OF HUMAN GENETICS, UNIVERSITY OF CHICAGO PRESS, CHICAGO,, US, vol. 45, no. 6, 1 December 1989 (1989-12-01), pages 889-904, XP000570035 ISSN: 0002-9297 ---	
P,X	DATABASE EM_HUM [Online] EMBL; 29 September 2000 (2000-09-29) SUGANO ET AL.: "Homo sapiens cDNA: FLJ21934 fis, clone HEP04364" retrieved from EBI, accession no. AK025587 Database accession no. AK025587 XP002186103 the whole document	3,11,12
P,X	-& DATABASE SWALL [Online] 1 March 2001 (2001-03-01) KAWABATA ET AL.: "cDNA: FLJ21934 fis, clone HEP04364" retrieved from EBI, accession no. Q9H6S4 Database accession no. Q9H6S4 XP002186764 the whole document -----	1

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 01/11869

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.: 20, 21, 23, 24  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-44 all partially, 45, 55 completely

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-44 all partially and 45 and 55 completely

An isolated polypeptide, an isolated polynucleotide and their uses where the polypeptide and polynucleotide are Seq Id Nos 1 and 11, respectively.

2. Claims: 1-44 all partially and 46 and 56 completely

Idem as for invention 1 above but for Seq Id Nos 2 and 12.

3. Claims: 1-44 all partially and 47 and 57 completely

Idem as for invention 1 above but for Seq Id Nos 3 and 13.

4. Claims: 1-44 all partially and 48 and 58 completely

Idem as for invention 1 above but for Seq Id Nos 4 and 14.

5. Claims: 1-44 all partially and 49 and 59 completely

Idem as for invention 1 above but for Seq Id Nos 5 and 15.

6. Claims: 1-44 all partially and 50 and 60 completely

Idem as for invention 1 above but for Seq Id Nos 6 and 16.

7. Claims: 1-44 all partially and 51 and 61 completely

Idem as for invention 1 above but for Seq Id Nos 7 and 17.

8. Claims: 1-44 all partially and 52 and 62 completely

Idem as for invention 1 above but for Seq Id Nos 8 and 18.

9. Claims: 1-44 all partially and 53 and 63 completely

Idem as for invention 1 above but for Seq Id Nos 9 and 19.

10. Claims: 1-44 all partially and 54 and 64 completely

Idem as for invention 1 above but for Seq Id Nos 10 and 20.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 18,21 and 24 are directed to methods of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

-----

Continuation of Box I.1

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

-----

Continuation of Box I.2

Claims Nos.: 20, 21, 23, 24

Claims 20,21,23,24 refers to agonists/antagonists of the polypeptide of claim 1 identified by the methods of claims 19 and 22 respectively, without giving a true technical characterization. Moreover, no such compounds are specifically defined in the description. It is only indicated that they "may include proteins" (such as antibodies)", nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of DME either by directly interacting with DME or by acting on components of the biological pathway in which DME participates." (page 31 lines 22-25; page 33 lines 23-27). In consequence the scope of said claims is ambiguous and vague, and their subject-matter is not sufficiently disclosed and supported ( Art. 5 and 6 PCT). No search can be carried out for such claims whose wording is, in fact, a mere recitation of the results to be achieved.

Claim 8 has only been search in as far as it concerns non-human transgenic organisms.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/11869

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9744466	A	27-11-1997	AU 2687897 A	09-12-1997
			WO 9744466 A1	27-11-1997
			EP 0904382 A1	31-03-1999
			JP 2000511046 T	29-08-2000
			US 6287834 B1	11-09-2001
-----				
EP 0905232	A	31-03-1999	EP 0905232 A1	31-03-1999
			US 2001024814 A1	27-09-2001
			CN 1214732 A	21-04-1999
			WO 9826053 A1	18-06-1998
-----				
EP 0463395	A	02-01-1992	EP 0463395 A1	02-01-1992
			AU 645091 B2	06-01-1994
			AU 7848791 A	02-01-1992
			CA 2045012 A1	23-12-1991
			JP 5211895 A	24-08-1993
			NZ 238610 A	25-11-1993
			US 5648482 A	15-07-1997
			US 5844108 A	01-12-1998
			DE 69126064 D1	19-06-1997
			DE 69126064 T2	28-08-1997
			ES 2103756 T3	01-10-1997
-----				

**THIS PAGE BLANK (USPTO)**

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant:

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**